arXiv:1501.04652v3 [math.QA] 4 Oct 2016

INTEGRATING QUANTUM GROUPS OVER SURFACES

DAVID BEN-ZVI, ADRIEN BROCHIER, AND DAVID JORDAN

Abstract. Braided tensor categories give rise to 4-dimensional topological field theories, extending constructions of Crane-Yetter-Kauffman in the case of modular categories. In this paper we apply the mechanism of factorization homology in order to construct and compute the category-valued invariants of surfaces which form the two-dimensional part of the theory. Starting from modules for the Drinfeld-Jimbo quantum group $U_q(\mathfrak{g})$ we obtain in this way a form of topologically twisted 4-dimensional $\mathcal{N} = 4$ super Yang-Mills theory, the setting introduced by Kapustin-Witten for the geometric Langlands program.

For punctured surfaces, in particular, we produce explicit categories which quantize character varieties (moduli of G-local systems) on the surface; these give uniform constructions of a variety of well-known algebras in quantum group theory. From the annulus, we recover the reflection equation algebra associated to $U_q(\mathfrak{g})$, and from the punctured torus we recover the algebra of quantum differential operators associated to $U_q(\mathfrak{g})$. From an arbitrary surface we recover Alekseev's moduli algebras. Our construction gives an intrinsically topological explanation for well-known mapping class group symmetries and braid group actions associated to these algebras, in particular the elliptic modular symmetry (difference Fourier transform) of quantum $\mathcal{D}$-modules.

Contents

1. Introduction 2
1.1. Factorization homology of surfaces 3
1.2. Summary of results 4
1.3. Detailed overview 6
1.4. Outlook 9
1.5. Acknowledgments 11
2. Factorization homology 12
2.1. Basic definitions 12
2.2. Pointed Structure 14
3. Categorical settings 14
3.1. An equivalence between **Rex** and **Pr** 15
3.2. The Deligne-Kelly tensor product 16
3.3. Tensor and braided tensor categories 17
4. Barr-Beck reconstruction of module categories 20
4.1. Barr-Beck-Tannaka-Krein reconstruction for braided tensor categories 23
5. Computing factorization homology of punctured surfaces 24
5.1. The distinguished object 24
5.2. Moduli algebras 25
5.3. Examples 28
5.4. Mapping class group actions 31
5.5. Braid group representations 32

6. Character varieties and quantization 32
6.1. Classical character varieties from Rep G 33
6.2. Gluing patterns, fat graphs and Poisson structures 33
6.3. Quantization via $\mathrm{Rep}_{\hbar} G$ 34
References 35

1. INTRODUCTION

This paper is concerned with interactions between braided tensor categories and topological field theory. One well-studied source of such interactions is provided by the identification of *modular* categories – braided tensor categories with strong finiteness – with the possible values of extended (3,2,1)-dimensional topological field theories on the circle. This is the Witten-Reshetikhin-Turaev (WRT) construction, originating in Chern-Simons theory.

Our interest is in a different interaction, tying braided tensor categories to *four*-dimensional topological field theories such as the topologically twisted $\mathcal{N} = 4$ super Yang-Mills theory used by Kapustin and Witten to study the geometric Langlands correspondence. In this story braided tensor categories appear as the possible values of a fully extended four-dimensional field theory on a point. More precisely, one considers the Morita theory of braided tensor categories; this forms a 4-category with appropriate bimodules as morphisms. The Cobordism Hypothesis [Lur] then allows us to build up towards a four-dimensional field theory, to an extent determined by finiteness properties enjoyed by the input.

The simplest example of such a four-dimensional theory is the four-dimensional "anomaly theory" for the WRT theory, which was first introduced by Crane-Yetter and Kauffman [CKY97] (and hence will be here-after abbreviated CYK), using a modular category construction. The CYK theory is revisited in recent work of Freed and Teleman [FT12], who show that a modular tensor category $\mathcal{M}$ defines a fully extended (in fact invertible) 4-dimensional topological field theory valued in the Morita 4-category of braided tensor categories.

We will be interested instead in braided tensor categories, such as the ribbon category $\mathcal{A} = \mathrm{Rep_q}\, G$ of representations of the quantum group, which are not necessarily finite or semi-simple, and hence not modular. Here $q \in \mathbb{C}$ is an arbitrary complex parameter: when q is a root of unity, $\mathcal{A}$ has the modular tensor category appearing in WRT and CYK theories as a semi-simple sub quotient. It is expected by experts (cf. [Wal]) that under mild conditions ribbon categories define "(3+1)-dimensional TFTs", meaning that they have all the lower-dimensional structures of a four-dimensional theory but are not defined on 4-manifolds (as follows for example from the infinite dimensionality of the vector spaces such a theory should attach to certain 3-manifolds).

In this paper we develop factorization homology of categories and apply it to explicitly construct and compute these topological field theories in dimensions at most two. In the case of $\mathrm{Rep_q}\, G$, we obtain categories which we call *quantum character varieties.* These quantize moduli spaces of local systems on surfaces and provide a unifying perspective on various constructions in quantum group theory. Quantum character varieties form the spectral side of the quantum Betti geometric Langlands conjecture [BZN16], and so should carry many rich structures which we plan to explore in future papers.

1.1. **Factorization homology of surfaces.** Factorization homology was originally introduced by Beilinson and Drinfeld [BD04] in the setting of conformal field theory, as an abstraction (and geometric interpretation) of the functor of conformal blocks of a vertex algebra. Factorization homology in the topological, rather than conformal, setting is developed in [Lur] and further in [AF12, AFT14], (see [Gin14] for a survey and [CG] for more general applications to quantum field theory).

The algebraic input to factorization homology of surfaces is an "E_2-algebra" – a.k.a. algebra over the little 2-disks operad – in an appropriate symmetric monoidal higher category $\mathcal{C}^{\boxtimes}$. Informally speaking, an E_2-algebra in $\mathcal{C}$ is an object $\mathcal{A} \in \mathcal{C}$ equipped with operations $\mathcal{A}^{\boxtimes k} \to \mathcal{A}$ parametrized in a locally constant fashion by framed embeddings of disjoint unions of k disks into a large disk, and satisfying a composition law governed by composition of disk embeddings. If we consider oriented, rather than only framed, embeddings of disks we obtain the notion of a *2-disk algebra*[1].

Our approach to constructing quantum character varieties is to apply the mechanism of factorization homology in the nonlinear setting of categories rather than the linear setting of vector spaces or chain complexes, i.e. we work in a 2-category $\mathcal{C}^{\boxtimes}$ of k-linear categories.[2]. An elementary but important observation is that in this case, an E_2-structure is determined already by the binary product $\mathcal{A}^{\boxtimes 2} \to \mathcal{A}$ labeled by a fixed embedding, an associator natural isomorphism, and a collection of braid group actions, given by monodromy over the configuration space. Taking into account compositions and coherences, one finds that an E_2-algebra in categories is simply a braided tensor category. Likewise, a 2-disk algebra in categories is a *balanced* (braided) tensor category (e.g. a pivotal braided category; see Remark 3.8).

The factorization homology of a framed surface S with coefficients in an E_2-algebra $\mathcal{A}$ is another object of $\mathcal{C}^{\boxtimes}$, denoted $\int_S \mathcal{A}$. The assignment,

$$S \longmapsto \int_S \mathcal{A} \in \mathcal{C}^{\boxtimes},$$

defines an invariant of framed manifolds, and is functorial for framed embeddings. When $\mathcal{A}$ is a 2-disk algebra, the assignment descends to an invariant of arbitrary *oriented* surfaces, and is functorial for oriented embeddings.

The invariant thus produced may be characterized as follows: to an open disk D^2 is assigned, by definition, the underlying object of $\mathcal{A}$; to a general surface S, is assigned the "integration" (i.e. co-limit) over all possible embeddings of a finite disjoint union of disks $i : (D^2)^{\sqcup k} \hookrightarrow S$. By construction the factorization homology of any surface S with coefficients in $\mathcal{A}$ carries a universal morphism of "localization",

$$\Delta_i : \mathcal{A}^{\boxtimes k} \to \int_S \mathcal{A},$$

for every disk embedding, which moreover factors through the E_2-multiplication (in our case, braided tensor structure), whenever it factors through a larger disk embedding. Finally, the unit of $\mathcal{A}$ endows the factorization homology of any surface with a canonical pointing: there is a distinguished object

---

[1] We caution that 2-disk algebras are historically called "framed" E_2-algebras, a potentially confusing terminology in the present context!

[2] Let us delay until Section 3 a discussion of the precise 2-categorical framework.

$$\mathcal{O}_{\mathcal{A},S} \in \int_S \mathcal{A}$$

which can be realized as

$$\mathcal{O}_{\mathcal{A},S} \simeq \Delta_i(1_{\mathcal{A}}),$$

for any disk embedding.

As with any homology theory, a primary tool for computing factorization homology is its *excision* property. Given an oriented 1-manifold M (in particular $M = S^1$ or $M = I$), the factorization homology $\int_{M\times\mathbb{R}} \mathcal{A}$ carries a canonical E_1-structure (i.e., an associative product; in our categorical setting, a tensor product) from the inclusion of disjoint unions of intervals inside a larger interval (i.e., we stack cylinders inside a larger cylinder). Moreover the invariant of a manifold with a collared boundary M is naturally a module for the E_1-structure on the invariant of its collared boundary. This structure gives rise the *excision* property of factorization homology: let $S = X^- \underset{M\times\mathbb{R}}{\sqcup} X^+$ be a collar gluing of $X^\pm$ along a 1-manifold P with a trivialization $N \cong M \times \mathbb{R}$ of a tubular neighborhood N of M. Then $\int_{X^\pm} \mathcal{A}$ are left and right $\int_{M\times\mathbb{R}} \mathcal{A}$-modules respectively, and there is an equivalence [AFT14, AF12] of categories

$$\int_S \mathcal{A} \simeq \int_{X^-} \mathcal{A} \underset{\int_{M\times\mathbb{R}} \mathcal{A}}{\boxtimes} \int_{X^+} \mathcal{A}.$$

1.2. **Summary of results.** Fix a balanced tensor category $\mathcal{A}$. Our main results are as follows:

- We construct the category $\int_S \mathcal{A}$ for any oriented surface S, equipped with a distinguished object $\mathcal{O}_{\mathcal{A},S}$ and carrying an action of the diffeomorphism group of S.
- To a surface with a marked S^1 boundary component we attach a canonical algebra object

  $$A_S := \underline{\mathrm{End}}_{\mathcal{A}}(\mathcal{O}_{\mathcal{A},S}) \in \mathcal{A},$$

  the internal endomorphism algebra of $\mathcal{O}_{\mathcal{A},S}$. In Theorem 5.11 we produce an equivalence,

  $$\int_S \mathcal{A} \simeq A_S\text{-mod}_{\mathcal{A}},$$

  as a module category for $\mathcal{A}$.
- In Section 5, we develop a combinatorial framework which allows us to produce explicit presentations of the algebras A_S from a combinatorial presentation ("handle and comb decomposition") of a punctured surface S. This involves giving an explicit computation of relative tensor products dictated by the excision axiom.

While most of our results apply to general braided tensor categories, we will mostly be concerned in this paper with the balanced tensor category $\mathcal{A} = \mathrm{Rep}_{\mathrm{q}}\, G$ of representations of the quantum group $U_q\mathfrak{g}$, associated to a reductive group G and an arbitrary $q \in \mathbb{C}^\times$ (see Section 1.3.2 for a precise definition). In the special case of the symmetric tensor category $\mathcal{A} = \mathrm{Rep}\, G$, the factorization homology $\int_S \mathrm{Rep}\, G$ makes sense on *any* topological space S. In the derived setting, it was proved in [BZFN10] that the result is the dg category of quasi-coherent sheaves on the derived *character stack* $\underline{\mathrm{Ch}}_G(S)$, the moduli stack of G-local systems on S (whose underlying underived stack parametrizes homomorphisms $\rho : \pi_1(S) \to G$, modulo conjugation in G):

$$\int_S \mathrm{Rep}_{dg}\, G \simeq QC_{dg}(\underline{\mathrm{Ch}}_G(S)).$$

The analogous result for abelian categories of quasi-coherent sheaves on character stacks of surfaces follows from the main results of this paper (see Remark 1.1 for the relation between the derived and abelian theories). Thus we will consider the category $\int_S \mathrm{Rep_q}\, G$ to be the quantum analog of sheaves on the character stack, and refer to it as the (G-)*quantum character variety*.

- For any surface the invariant part of the algebra A_S is an explicit modular invariant quantization of the Poisson algebra of functions on the G-character variety $\mathrm{Ch}_G(S)$ (Section 6).
- For the annulus A_S recovers the reflection equation algebra $\mathcal{O}_q(G)$. For the punctured torus A_S recovers the algebra $\mathcal{D}_q(G)$ of quantum differential operators on G. More generally, for a punctured surface of arbitrary genus we can recover so-called moduli algebras of Alekseev.

1.2.1. *Factorization homology of linear categories.* In order to set up factorization homology of linear categories, one must find a suitable higher categorical framework — specifically we require a higher category of categories which carries a symmetric monoidal structures and is closed under (sufficiently many) colimits. The collection of abelian categories fails to satisfy these properties, and must be enlarged, in one of several related ways. In setting up the general foundations we establish a formalism that is potentially of independent interest to specialists in the theory of tensor categories.

- We define factorization homology of tensor categories in the setting **Pr** of compactly generated presentable (or equivalently finitely cocomplete **Rex**) categories.
- We develop several techniques, related to Beck monadicity, for describing module categories explicitly as categories of modules for algebra objects, mostly extending well-known results from the setting of finite (and often semi-simple) tensor categories (cf. [ENO10, Ost03, DSPS13]) to the infinite setting. In particular we prove monadicity results describing various constructions with abelian tensor categories (such as base changes, relative tensor products, and traces) as categories of modules over internal endomorphism algebras.

Remark 1.1 (Derived version)**.** Prompted by discussions following the posting of our preprint, J. Lurie [Lur16] has proved that (contrary to widely held expectations) the collection of Grothendieck abelian categories is closed under tensor products. The result is part of a general theory of derived analogues, called Grothendieck prestable ∞-categories, which are roughly (the positive halves of) differential graded categories with t-structures whose heart is a Grothendieck abelian category. It follows from Lurie's results that one can define factorization homology of E_2-categories either in the setting of dg (or stable ∞) categories, or in the refined setting of Grothendieck prestable categories, i.e., keeping track of t-structures. For rigid braided tensor categories one can check that our constructions in the discrete setting are compatible with their derived analogues – i.e., our quantum character varieties are the hearts of the natural derived quantum character varieties, obtained by integrating the dg categories of representations of quantum groups and keeping track of t-structures. (We intend to return to this derived comparison in a later paper.) In particular this implies the categories constructed in this paper are in fact abelian – a fact which is evident for the categories on punctured surfaces we study in detail.

Remark 1.2 (Skein Categories)**.** It is interesting to compare factorization homology with the theory of skein algebras (see e.g. [RS02, FG00]). Skein algebras provide a convenient graphical calculus for quantizing SL_2 (and more generally SL_n) character varieties and constructing associated 3-manifold invariants. Our factorization homology construction can be viewed as a general formalism extending skein theory to arbitrary braided tensor categories. Roughly speaking, an object of a skein category associated to a surface S, a braided tensor category $\mathcal{A}$ and a presentation of $\mathcal{A}$ is a configuration of disks on S, each disk colored with a generator of $\mathcal{A}$. Homomorphisms between such objects are defined by quotienting vector spaces of colored tangles, according to "local relations" in $\mathcal{A}$; the result is not typically an abelian category. There is an evident functor to the corresponding factorization homology category, which provides an abelian co-completion. Factorization homology also enjoys manifest locality and functoriality properties, independence of presentation, good behavior for closed as well as open surfaces and a natural (and nontrivial) derived extension, which make it readily applicable to problems in geometric representation theory. It would be very interesting to identify our factorization homology construction and the similarly general abstract skein theory developed by Walker [Wal].

1.3. **Detailed overview.** We now describe our main results in more detail.

1.3.1. *Punctured surfaces.* Our main result, Theorem 5.11, is a concrete computation of the quantum character variety of an arbitrary punctured surface S. Recall that the inclusion of a disk around a point $x \in S$ in a surface defines a localization functor

$$\Delta_x : \mathcal{A} \to \int_S \mathcal{A}, \qquad \mathbf{1}_{\mathcal{A}} \mapsto \mathcal{O}_{\mathcal{A},S}$$

from $\mathcal{A}$ to the quantum character variety, sending the unit $\mathbf{1}_{\mathcal{A}}$ to the "quantum structure sheaf", the distinguished object $\mathcal{O}_{\mathcal{A},S} \in \int_S \mathcal{A}$. In the commutative case $\mathcal{A} = \operatorname{Rep} G$ localization is given by pulling back sheaves under

$$\underline{\mathrm{Ch}}_G(S) \longrightarrow \underline{\mathrm{Ch}}_G(D^2) \simeq pt/G,$$

and the distinguished object is the structure sheaf $\mathcal{O}_{\underline{\mathrm{Ch}}_G(S)}$ of the classical character stack. Unlike the commutative case, however, the quantum character variety does not carry an $\mathcal{A}$-module structure in general. This is the two-dimensional analog of the assertion that Hochschild homology $\int_{S^1} A$ of an associative algebra A carries a trace map from A but has no natural A-module structure.

Our description of quantum character varieties is based on the observation that if we chose a boundary point $x \in \partial S$ we do obtain a natural $\mathcal{A}$-module structure on $\int_S \mathcal{A}$. In fact this is naturally a braided module category; this is developed and exploited in [BZBJ16]. We can then describe the entire category $\int_S \mathcal{A}$ as the category of modules in $\mathcal{A}$ for the internal endomorphism algebra of the distinguished object, relative to this $\mathcal{A}$-module structure:

Theorem 1.3 (Theorem 5.11)**.** *Let* $A_S := \underline{\mathrm{End}}_{\mathcal{A}}(\mathcal{O}_{\mathcal{A},S})$ *denote the internal endomorphism algebra of* $\mathcal{O}_{\mathcal{A},S}$*. We have an equivalence of categories*

$$\int_S \mathcal{A} \simeq A_S\text{-mod}_{\mathcal{A}}$$

respecting natural actions of the mapping class group of S *relative to the boundary.*

Remark 1.4. For $\mathcal{A} = \mathrm{Rep}\ G$, this captures the statement that $\underline{\mathrm{Ch}}_G(S)$ is affine over pt/G.

The algebra A_S can be described in completely explicit terms, once one chooses a topological presentation of the surface. Given some additional combinatorial data, which we call a "gluing pattern" P for S (See Figure 1), we define an algebra a_P in $\mathcal{A}$ via a "generators and relations" presentation.

Theorem 1.5. *Given a gluing pattern* P *for* S *there is a canonical isomorphism* $A_S \cong a_P$ *of algebras in* $\mathcal{A}$*, where*

$$a_P \simeq (\mathcal{O}(\mathcal{A}))^{\otimes 2g+r-1}$$

where each tensor factor is a sub-algebra, and where cross relations are expressed using the braiding on $\mathcal{A}$*, according to the data of the gluing pattern* P*.*

The proof is based on applications of the Barr-Beck monadicity theorem developed in Section 3, which allow us to the describe various module categories over a tensor category $\mathcal{A}$ as categories of modules over an algebra internal to $\mathcal{A}$.

Geometrically, this explicit presentation is related to the fact that the fundamental group of a punctured surface is a free group of rank $2g + r - 1$. The gluing pattern in particular highlights a set of free generators, around which the presentation for the algebra a_P is organized. We emphasize, however, that in general a_P is *not* simply the braided tensor product of copies of the $\mathcal{O}(\mathcal{A})$'s – this appears only in the case of the many-punctured disk – but rather the relations depend in an interesting way on the pattern P: see Section 5.

The disk, annulus and once-punctured torus each admit a unique gluing pattern; hence we will denote these simply by D^2, Ann, $T^2\backslash D^2$, respectively. For higher genus and number of punctures, a given surface may admit several distinct gluing patterns; each such gives a different presentation for the algebra A_S.

1.3.2. *The case of quantum groups.* Our main example will come from fixing a reductive algebraic group G, and a Killing form κ on $\mathfrak{g} = Lie(G)$. We will consider the balanced tensor category $\mathcal{A} = \mathrm{Rep_q}\, G$: by this notation we will denote either the category of finite dimensional $U_q(\mathfrak{g})$-modules, where $U_q(\mathfrak{g})$ is the quantum group associated to $\mathfrak{g}$ and κ when G is simply connected, or more generally the finite-index braided tensor subcategory of $U_q(\mathfrak{g})$ corresponding to G (and determined by its Cartan subgroup) when G is not simply connected. We do not recall the a presentation of $U_q(\mathfrak{g})$ here; it can be found e.g. in [CP94, Section 9.1]). When q is a root of unity, there are several different versions of $\mathrm{Rep_q}\, G$ one may consider, and to which our results apply; see Section 1.4.3). We also make use of the ind-completion of $\mathrm{Rep_q}\, G$, consisting of locally finite-dimensional $U_q(\mathfrak{g})$-modules.

When $\mathcal{A} = \mathrm{Rep_q}\, G$, the algebras a_P recover several interesting and well-known constructions in the geometric representation theory of quantum groups:

(1) For the disk, we have $a_{D^2} \cong \mathbf{1}_{\mathcal{A}}$, which is just the tautological equivalence $\mathcal{A} \simeq \mathbf{1}_{\mathcal{A}}\text{-mod}_{\mathcal{A}}$.

(2) For the annulus or cylinder, we have $a_{Ann} = \mathcal{O}(\mathcal{A}) \cong \mathcal{O}_q(G)$, the reflection equation algebra [DKM03, DM03, DM02b, DM02a].

(3) For the punctured torus, we have $a_{T^2\backslash D^2} \cong \mathcal{D}_q(G)$, the algebra of quantum differential operators on G. This algebra has received a lot of attention in recent years [STS94, Ale93, AGS96, BK08, BK06, VV10, Jor09, Jor14, BJ14].

(4) More generally, the moduli algebras of Alekseev [Ale93], [AGS96] can be recovered as algebras a_P associated to certain gluing patterns for higher genus surfaces.

In particular, it should be emphasized that the algebras a_P have explicit generators-and-relations presentations, PBW bases, and interesting representation theory related to the symplectic geometry of the classical character variety. These features are not typical of factorization homology in general, but emerge from the representation theoretic framework.

On the topological side, the reflection equation algebra, quantum differential operator algebra, and moduli algebras all admit interesting actions of mapping class groups of their associated surfaces, and can moreover be used to produce representations as in of the associated surface braid groups [Jor09, BJ14], extending Reshetikhin and Turaev's constructions [RT90]. Whereas these topological structures have historically been themselves constructed via generators-and-relations comparison, they now follow naturally from the topological framework of factorization homology. This is an important distinction if one wants to produce topological invariants and categorical structures from these constructions - the categories and distinguished objects associated to surfaces are functorial, but the particular presentations are not.

1.3.3. *Quantization of character varieties.* Let S be a surface. We let $\mathfrak{R}_G(S)$ denote the affine variety

$$\mathfrak{R}_G(S) = \{\rho : \pi_1(S) \longrightarrow G\}$$

of homomorphisms from the fundamental group of S to the group G. Equivalently, $\mathfrak{R}_G(S)$ is the moduli space of local systems equipped with a trivialization at a fixed point. Thus sheaves on $\underline{\mathrm{Ch}}_G(S)$ are given by modules over the corresponding algebra of global functions as an object in $QC(pt/G) = \mathrm{Rep}\, G$, while sheaves on $\mathfrak{R}_G(S)$ are recovered by de-equivariantization. We denote by $\mathrm{Ch}_G(S)$ the G-character *variety* (not stack), that is the affine categorical quotient

$$\mathrm{Ch}_G(S) := \mathfrak{R}_G(S) /\!\!/ G$$

by the natural adjoint action of G, to distinguish it from the character stack $\underline{\mathrm{Ch}}_G(S)$. The space $\mathrm{Ch}_G(S)$ carries a canonical Poisson structure originally due to Atiyah–Bott [AB83] and Goldman [Gol84]. A discrete, combinatorial construction of this structure was given by Fock–Rosly [FR99] using classical r-matrices. They construct a Poisson structure on the representation variety $\mathfrak{R}_G(S)$ itself, which depends on the choice of a representation of S as a so-called ciliated graph. When $\mathcal{A}$ is the category of modules over the formal quantum group $U_\hbar(\mathfrak{g})$ and P is a gluing pattern for S, the algebra a_P is a flat deformation of the algebra of functions on the representation variety $\mathfrak{R}_G(S)$. By regarding a gluing pattern P for S as a ciliated graph with only one vertex, the Fock-Rosly construction determines a Poisson structure on $\mathfrak{R}_G(S)$.

Fock–Rosly's construction was partly inspired by the work of Semenov-Tian-Shansky [STS94] who introduced a certain dual Poisson structure on G, characterized by a classical version of the reflection equation, and a Poisson structure on $G \times G$ (the classical Heisenberg double) thought as a Poisson-Lie version of the cotangent bundle T^*G. He also constructed quantization of those structures. The relation between those Poisson structures and $\mathrm{Ch}_G(S)$ was already noticed in [AM95].

Quantizations of $\mathrm{Ch}_G(S)$ were then obtained in [Ale93, AGS96] by, roughly, replacing classical r-matrices by quantum R-matrices, hence giving an FRT-like (in the sense of [FRT90]) presentation of the sought quantization. We prove the following:

Theorem 1.6. *The algebra a_P is a quantization of the Fock–Rosly Poisson structure on $\mathfrak{R}_G(S)$ associated with P. Its $U_\hbar(\mathfrak{g})$-invariant part is independent of P, and is a quantization of the canonical Poisson structure on $\mathrm{Ch}_G(S)$.*

For suitable choices of gluing pattern P, our quantizations recover Alekseev's algebras. Actions of the mapping class group of the underlying surface on quantizations of character varieties are constructed in [AS96] directly via generators and relations. In our approach this is rather a by-product of their topological definition via factorization homology.

1.4. **Outlook.** Here we collect several remarks pertaining to further directions of study springing from the current work.

1.4.1. *Computations in closed surfaces.* In the tandem paper [BZBJ16], we extend the present techniques to the setting of closed, and possibly marked, surfaces. The key technical difficulty there is that, while punctured surfaces can be glued from disks along boundary intervals, closed surfaces require gluing along boundary annuli as well. In particular, in order to understand the algebraic data involved in sealing up punctures, or gluing in marked points, we need a convenient monadic framework for working with module categories over the factorization homology, $\int_{Ann}\mathcal{A}$, of the annulus. This is accomplished via the theory of quantum moment maps and quantum Hamiltonian reduction.

1.4.2. *Four-dimensional topological field theory.* Factorization homology fits naturally into the language of extended topological field theory and the cobordism hypothesis [Lur09, CS]. This connection is motivational, but not technically necessary for our paper, so we will be informal in our treatment.

The Cobordism Hypothesis proved by Lurie [Lur09] establishes that fully extended n-dimensional topological field theories are functorially determined by the invariant assigned to an n-disk, an object of a higher category with suitable finiteness conditions. An important and accessible special case is provided by the theory[3] of E_n-algebras, i.e., algebras over the little n-disks operad, which automatically satisfy the finiteness conditions necessary to define invariants of manifolds of dimension at most n [Lur09, CS]. In this setting the invariants are given by factorization homology.

The factorization homology of a balanced tensor category $\mathcal{A}$ over surfaces forms the (0,1,2)-dimensional part of a 4-dimensional oriented topological field theory – equivalently, a category-valued 2-dimensional topological field theory. To a closed surface we attach the category $\int_S\mathcal{A}$. To a 1-manifold M we attach the 2-category $\int_{M\times\mathbb{R}}\mathcal{A}$-mod. Thus for a surface with collared boundary we have an object in the 2-category attached to the boundary:

$$\int_S \mathcal{A} \in \left(\int_{\partial S\times\mathbb{R}} \mathcal{A}\right)\text{-mod}$$

---

[3] More precisely, the *Morita* theory, where E_n-algebras in a k-category $\mathcal{C}^{\boxtimes}$ form an $n{+}k$-category, generalizing the familiar Morita theory of associative algebras for $n = 1$.

and the sewing property above provides the composition structure for the topological field theory. More generally we obtain invariants for surfaces with boundary components marked by module categories for $\int_{S^1\times\mathbb{R}}\mathcal{A}$. Finally to a point we attach the 3-category $\mathcal{A}$-mod-mod of $\mathcal{A}$-linear two-categories (module categories for the monoidal 2-category of $\mathcal{A}$-module categories), which corresponds to $\mathcal{A}$ under the natural embedding of the Morita theory of braided categories into the 4-category of 3-categories.

While the computations in the present paper and [BZBJ16] are restricted to dimension 2, it is expected that the topological field theory defined by $\mathcal{A}$ extends to dimension three. This involves showing that a rigid braided tensor category $\mathcal{A}$ defines a 3-dualizable object in the Morita 4-category of E_2-algebras in **Rex** (see [JFS15], [Hau14] for constructions of higher Morita categories), and that a ribbon structure gives rise to a homotopy fixed point structure for the induced $SO(3)$-action. This is the subject of work in progress with N. Snyder.

1.4.3. *Roots of unity.* The techniques developed in this paper apply to arbitrary braided tensor categories. In particular, when we consider $\mathrm{Rep_q}\,G$, for q a root of unity, there are four distinct settings in which we can work. The De Concini-Kac quantum group $U_q^{DK}(\mathfrak{g})$ arises from directly specializing q in Serre's presentation for the quantum group [DCK90]. The resulting algebra has a large center, over which it decomposes (étale-locally) as a direct sum of matrix algebras. The small quantum group, $u_q\mathfrak{g}$ appears as a quotient of $U_q^{DK}(\mathfrak{g})$ at a certain central character [Lus90]; while $u_q(\mathfrak{g})$ does not admit an R-matrix [LN15], certain finite-degree extensions of it do. There is Lusztig's restricted quantum group $U_q^{res}(\mathfrak{g})$, which includes divided powers of Serre generators E_i and F_i, whose category of representations give a braided tensor category. Finally, there is the braided tensor subcategory of tilting modules [APK91] for $U_q^{res}(\mathfrak{g})$, and its quotient by negligible morphisms, the modular tensor category $\mathcal{M}_q$ of Reshetikhin-Turaev theory [RT91]. The four-dimensional TFT defined by $\mathcal{M}_q$ has already been studied by Crane-Kauffman-Yetter [CKY97], but the others appear not to have received the same attention. It should be very interesting to compare the constructions in the present paper in each of the different root of unity settings outlined above.

1.4.4. *Betti Geometric Langlands.* An important motivation for this series of papers is the formulation of a Betti form of the quantum geometric Langlands conjecture. The spectral side of the Betti Geometric Langlands Conjecture [BZN16] replaces coherent sheaves on the de Rham space of flat G-connections on an algebraic curve X [AG14] by coherent sheaves on the Betti space, the character variety $\underline{\mathrm{Ch}}_G(S)$ of the underlying topological surface. The quantization of the de Rham category is constructed using representations of affine Kac-Moody algebras – in particular through the localization functor studied by Beilinson-Drinfeld from LG_+-integrable $\mathfrak{g}$-modules to twisted $\mathcal{D}$-modules on $Bun_G(X)$ (depending on a point $x \in X$). Likewise the quantization of the Betti categories is constructed in this paper by assembling the localization functors Δ_x $(x \in S)$ from representations of quantum groups. The quantum analog of the Riemann-Hilbert correspondence is provided by the Kazhdan-Lusztig equivalence between the two representation categories. We summarize the situation in the following diagram, where dotted arrows indicate equivalences after analytification:

$$\begin{tikzcd}
\operatorname{Rep_q} G \arrow[r, "\Delta_x"] \arrow[d, "KL"', "\sim"] & \int_S \operatorname{Rep_q} G \arrow[r, rightsquigarrow, "q\to 1"] \arrow[d, dotted, "q-RH"'] & \underline{\mathrm{Ch}}_G(S) \arrow[d, dotted, "RH"'] \\
(\widehat{\mathfrak{g}}\text{-mod}_k)^{LG_+} \arrow[r, "\Delta_x"] & \mathcal{D}_k(Bun_G(X)) \arrow[r, rightsquigarrow, "k\to\infty"] & Conn_G(X)
\end{tikzcd}$$

The quantum Betti conjecture [BZN16] relates the categories constructed in this paper with twisted sheaves with nilpotent singular support on $Bun_{G^\vee}(X)$. This conjecture is motivated in turn by the work of Kapustin-Witten [KW07], in which Langlands duality is related to electric-magnetic S-duality in 4-dimensional topological field theory ($\mathcal{N} = 4$ supersymmetric Yang-Mills theory in the GL twist). The topological field theory constructed in this paper using factorization homology is an algebraic model for this topological field theory (in contrast with the de Rham version which is not topological).

S-duality is also expected to have another analytic (though not algebraic) manifestation, as a duality between quantum character varieties for Langlands dual groups at dual levels (roughly $q = e^{\pi ik}$ and $q^\vee = e^{-\pi i/k}$). This is a manifestation of the celebrated but mysterious modular invariance in representation theory of quantum groups: certain aspects of the representation theory of $U_q\mathfrak{g}$ depend naturally not on q but only on the corresponding elliptic curve. This modularity is expressed by Faddeev's modular double [Fad00] and many subsequent works (see for example [TV14, FI14] and references therein) and the Langlands duality for quantized cluster varieties of [FG09].

1.4.5. *The KZ category.* The Poisson structure on character varieties can also be obtained by an appropriate reduction of a certain *quasi*-Poisson structure on $\mathfrak{R}_G(S)$ introduced in [AKSM02]. This construction was extended and somewhat simplified in [LBS13] and quantized by the same authors in [LBS14]. The quantization of such a quasi-Poisson structure is an algebra internal to the Drinfeld braided tensor category constructed from Rep G using a Drinfeld associator. While the introduction of associators makes the quantizations of [LBS14] somewhat less explicit than the moduli algebras of [Ale93], the role of braided tensor categories is made conceptually clearer. We note in particular that the "fusion" procedure of [LBS14] is very similar to the construction of a_P in Section 5, from copies of $\mathcal{O}(\mathcal{A})$.

We thus conjecture that their construction agrees with ours when $\mathcal{A}$ is the Drinfeld category. This would imply in particular that the algebras they obtain are related to those of Alekseev–Grosse–Schomerus by the Kohno–Drinfeld equivalence. We also expect the representations of surface braid groups obtained from factorization homology with coefficients in the Drinfeld category to coincide with those coming from the monodromy of the KZB equations [CEE09, Enr14]. This would give an explicit computation of the latter using quantum groups.

1.5. **Acknowledgments.** We would like to thank Jacob Lurie for first suggesting the $\operatorname{Rep_q} G$ TFT as a Betti setting for quantum geometric Langlands and for patiently answering our many questions on Grothendieck categories, Pavel Etingof and Benjamin Enriquez for sharing their ideas concerning elliptic structures on categories, and Dan Freed and Kevin Walker for sharing their understanding of topological field theory and braided tensor categories and their closely related works in progress with Teleman and Vazirani, respectively. We'd also like to thank Martin Brandenburg, Steve Lack, Mike Shulman, Theo Johnson-Freyd and Ross Street, and especially Daniel Schäppi and Chris Schommer-Pries for many helpful comments on 2-categorical issues; John Francis, Greg Ginot, Owen Gwilliam, Claudia Scheimbauer, and Noah Snyder for discussions on factorization homology, and Anatoly Preygel for a discussion of t-structures. We would like to thank Sam Gunningham, David Nadler, and Monica Vazirani for countless discussions surrounding geometric Langlands and related topics

The work of DBZ was partly supported by NSF grant DMS-1103525. The work of DJ was supported by NSF grant DMS-1103778 and ERC grant QuantGeomLangTFT 637618. DBZ and DJ would like to acknowledge that part of the work was carried out at MSRI as part of the program on Geometric Representation Theory.

2. Factorization homology

We review factorization homology following [AFT14, AF12] to which we refer for details; see also [Gin14] for a survey of the theory. In this paper we focus on factorization homology over compact surfaces with boundaries. However we will also need to compute factorization homology on simple examples of *manifolds with corners*, i.e. 2-dimensional (paracompact Hausdorff) topological spaces locally modeled on $\mathbb{R}^k_{\geq 0} \times \mathbb{R}^{2-k}$. Those are particular examples of stratified manifolds, on which factorization homology is well-defined thanks to [AFT14].

2.1. **Basic definitions.** Let Mfld^2_{fr} (resp. Mfld^2_{or}) be the $(\infty, 1)$ category whose objects are framed (resp. oriented) 2-dimensional manifolds with corners, and morphisms between manifolds M, N is the ∞-groupoid $Emb(M, N)$ of embeddings respecting the framing (resp. the orientation). Hence 1-morphisms are smooth embeddings, 2-morphisms are paths between embeddings (i.e. isotopies), 3-morphisms are homotopies between those and so on. The disjoint union turns those categories into symmetric monoidal categories. Let $\mathrm{Disk}^2_{\partial,fr}$ (resp. $\mathrm{Disk}^2_{\partial,or}$) denote the full subcategory generated under disjoint union by the disks $\mathbb{R}^k \times \mathbb{R}^{2-k}_{\geq 0}$, $0 \leq k \leq 2$ equipped with their standard framing (resp. orientation). We define similarly $Disk^{fr}_2$ and Disk^2_{or} as the categories generated by $\mathbb{R}^2$. Fix an $(\infty, 1)$-symmetric monoidal category $(\mathcal{C}, \boxtimes)$.

Definition 2.1. A $\mathrm{Disk}^2_{s,B}$-algebra in $\mathcal{C}$, for $s \in \{fr, or\}$, $B \in \{\emptyset, \partial\}$, is a symmetric monoidal functor from $\mathrm{Disk}^2_{s,B}$ to $\mathcal{C}$.

Remark 2.2. A Disk^2_{fr}-algebra (or rather the image of $\mathbb{R}^2$) is usually called an E_2-algebra, or algebra over the little disk operad. Similarly a Disk^2_{or}-algebra is an algebra over the framed little disk operad. A $\mathrm{Disk}^2_{fr,\partial}$-algebra is equivalent to the data of a triple of objects $(\mathcal{A}, \mathcal{M}, \mathcal{N})$, the image of the triple

$$(\mathbb{R}^2, \mathbb{R} \times \mathbb{R}_{\geq 0}, \mathbb{R}^2_{\geq 0}),$$

where $\mathcal{A}$ is an E_2-algebra and $\mathcal{M}, \mathcal{N}$ are $\mathcal{A}$-modules satisfying several conditions (e.g. the pair $(\mathcal{A}, \mathcal{M})$ is an algebra over the Swiss-Cheese operad [Vor99]). In this paper we focus to the particular case $\mathcal{M} = \mathcal{N} = \mathcal{A}$, i.e. all disks are sent to the same category regardless of their manifold with corner structure.

We now assume following [AF12], [Lur09] that $\mathcal{C}$ is cocomplete and that for every $c \in \mathcal{C}$, the functor $c \otimes -$ commutes with small colimits. In this setting the following colimit uniquely defines the factorization homology with coefficients in an E_2-algebra $\mathcal{A} \in \mathcal{C}$:

Definition 2.3 ([AF12, Lur])**.** Factorization homology with coefficients in an E_2 (resp. $Disk_2$) algebra $\mathcal{A} \in \mathcal{C}$ is the left Kan extension of the above symmetric monoidal functor with respect to the inclusion

$$\mathrm{Disk}^2_{s,\partial} \longrightarrow \mathrm{Mfld}^2_s$$

where $s \in \{fr, or\}$, which will be denoted by

$$M \longmapsto \int_M \mathcal{A}.$$

Let Y be a 1-dimensional manifold with corners. Set $Y^+ = Y \times \mathbb{R}_{\geq 0}$ and $Y^- = Y \times \mathbb{R}_{\leq 0}$. Then $Y^\pm$ are (framed, oriented) manifolds with corners. If $X^\pm$ are manifolds with corners and

$$\iota^\pm : Y^\pm \longrightarrow X^\pm$$

are framing (resp. orientation) preserving smooth embeddings such that

$$\iota^\pm(Y^\pm \times \{0\}) \subset \partial X^\pm,$$

then the gluing of $X^\pm$ along Y is a well defined (framed, oriented) manifold with corner.

One of the main properties of factorization homology is the following:

Theorem 2.4. *Let $\mathcal{A}$ be an E_2 (resp. framed E_2) algebra in $\mathcal{C}$. Then the functor $\int_{(-)} \mathcal{A}$ satisfies, and is characterized, by the following properties:*

- *If U is contractible, then there is an equivalence of categories*

$$\int_U \mathcal{A} \simeq \mathcal{A}.$$

- *A homeomorphism $U \cong P \times \mathbb{R}$, for a 1-dimensional manifold with corners P, equips $\int_U \mathcal{A}$ with a canonical monoidal (E_1) structure (from inclusions of intervals inside a larger interval).*
- *Let $U = X^- \underset{P \times \mathbb{R}}{\sqcup} X^+$ be a collar gluing of $X^\pm$ along a codimension 1 sub-manifold P with a choice of a trivialization $N \cong P \times \mathbb{R}$ of a tubular neighborhood N of P. Then $\int_{X^\pm} \mathcal{A}$ are left and right $\int_{Y \times \mathbb{R}} \mathcal{A}$-modules respectively, and there is an equivalence of categories*

$$\int_U \mathcal{A} \simeq \int_{X^-} \mathcal{A} \underset{\int_{Y \times \mathbb{R}} \mathcal{A}}{\boxtimes} \int_{X^+} \mathcal{A}.$$

Remark 2.5. Along the lines of Remark 2.2, in the particular case considered in this paper, $\int_M \mathcal{A}$ depends only on the manifold with boundary underlying M, not of its manifold with corners structure. This will allow us to see surfaces with circle boundaries as being obtained by gluing closed disks whose boundary is divided into several intervals, and then smoothing the remaining corners.

Remark 2.6. Let S be a surface with, say, a single, circular boundary, and let $\tilde{S}$ be the manifold with corners obtained by subdividing the boundary of S into n intervals. Then, providing that the framings match, or that $\mathcal{A}$ is balanced, $\int_{\tilde{S}} \mathcal{A}$ has a natural structure of a module over

$$\int_{\sqcup_n \mathbb{R}^2} \mathcal{A} \simeq \mathcal{A}^{\boxtimes n}.$$

On the other hand, $\int_S \mathcal{A}$ is naturally a $\int_{S^1\times\mathbb{R}} \mathcal{A}$-module. By the previous remark, we have an equivalence of categories

$$\int_S \mathcal{A} \simeq \int_{\tilde{S}} \mathcal{A},$$

hence the marking induces an $\mathcal{A}^{\boxtimes n}$-module on $\int_S \mathcal{A}$ as well. It is easily seen that this module structure is isomorphic to the one obtained via the composition

$$\mathcal{A}^{\boxtimes n} \longrightarrow \mathcal{A} \longrightarrow \int_{S^1\times\mathbb{R}} \mathcal{A}.$$

2.2. **Pointed Structure.** An important additional feature of factorization homology is that it is a *pointed* theory: the invariant assigned to an n-manifold M by an E_n-algebra comes equipped with a canonical morphism from the unit of the target category $\mathcal{C}$, constructed as follows. Let $\emptyset$ be the empty manifold. As $\emptyset$ is the unit for the disjoint union, we have $\int_\emptyset \mathcal{A} = \mathbf{1}_\mathcal{C}$. Moreover, $\emptyset$ is an initial object in $\mathrm{Mfld}^2_{or/fr}$; for any manifold M, there is a unique embedding:

$$\emptyset \longrightarrow M.$$

Its image through factorization homology provides a distinguished morphism

$$\mathbf{1}_\mathcal{C} \longrightarrow \int_M \mathcal{A}.$$

In the case that $\mathcal{C}$ is itself some collection of linear categories (as in the next section), we have that $\int_M \mathcal{A}$ is a category, $\mathbf{1}_\mathcal{C}$ is a category of vector spaces, so that this pointed structure produces a distinguished object in it, denoted $\mathcal{O}_{\mathcal{A},M}$, the image of the one-dimensional vector space. Distinguished objects play a key role in our main result, so we will detail some of their properties in Section 5.1.

3. CATEGORICAL SETTINGS

We now describe the categorical setting in which we will operate. It is important for constructions such as factorization homology to work in an ambient category (or ∞-category) with enough colimits and a symmetric monoidal structure preserving those colimits. We will largely work in the "discrete" setting of 1-categories, so that only a passing knowledge of ∞-categories is necessary to read the paper. However, we must address the existence of co-limits in various 2-categories of categories. We then recall the basic notions of tensor categories, braided tensor categories, module categories, etc.

It is known that the familiar setting **Ab** of small **k**-linear abelian categories does not possess these features, since the Deligne tensor product of abelian categories is no longer abelian in general [LF12]. In this section we will recall three convenient categorical settings where this framework is well-defined, and discuss the compatibility between them. We will consider the following 2-categories:

- **Rex**, of finitely co-complete **k**-linear categories with right exact functors,
- **PR**, of presentable **k**-linear categories with cocontinuous functors,
- **Pr**, of compactly generated presentable categories with compact and cocontinuous functors (also known as locally finitely presentable, $\mathbf{LFP_k}$).

We also mention the related setting of Grothendieck abelian categories (which are in particular presentable), see Remark 1.1. As motivating examples, categories of finite-dimensional (or alternatively, finitely presented) modules over a **k**-algebra, as well as finite dimensional representations of an algebraic group are **Rex**, categories of all modules over an **k**-algebra are **Pr** and Grothendieck. Likewise coherent sheaves on varieties or stacks are **Rex** while quasi-coherent sheaves are **Pr** and Grothendieck.

Terminology. By a *2-category* we will always mean a (non-strict) $(2,1)$-category, in other words, we will not use any non-invertible 2-morphisms. Equivalently, this means an $(\infty,1)$-category with 1-truncated mapping spaces. We suppress set theoretic issues — in particular all colimits will be small without further comment.

A category is **k**-*linear* if it is enriched over the category of **k**-vector spaces (note there are no finiteness assumptions on Homs or lengths of objects). We will always work with categories enriched over some field **k**, usually $\mathbf{k} = \mathbb{C}$, and without further comment except when technically necessarily.

3.1. **An equivalence between Rex and Pr.** Recall that a category is presentable (also known as locally presentable) if it is accessible (generated under colimits by a small subcategory) and cocomplete (closed under small colimits – in fact this implies it is complete as well). We denote by **PR** the 2-category of presentable **k**-linear categories with colimit preserving (aka cocontinuous) functors and natural isomorphisms.

Presentable categories provide a very flexible setting for algebra: tensor products of presentable categories are again presentable; so are colimits of presentable categories; the adjoint functor theorem provides right adjoints to colimit preserving functors, to which we can then apply techniques such as Barr-Beck monadicity to describe categories explicitly as categories of (co)modules. In practice the presentable categories we encounter on punctured surfaces will all be (Grothendieck) abelian categories. (See also Remark 1.1.)

Recall that an object $c \in \mathcal{C}$ is compact if $Hom(c,-)$ preserves filtered colimits, and a category $\mathcal{C}$ is said to be compactly generated if any object is a filtered colimit of compact objects (equivalently, we can write $\mathcal{C} = \operatorname{ind}\mathcal{C}_c$ as the ind-category of its full subcategory of compact objects). We denote by $\mathbf{Pr} \subset \mathbf{PR}$ the subcategory of compactly generated **k**-linear presentable categories and *compact* functors, i.e., functors preserving compact objects or equivalently (by the adjoint functor theorem) possessing right adjoints preserving filtered colimits.

A category is *finitely co-complete* if it admits finite colimits. We let **Rex** denote the 2-category of essentially small finitely cocomplete categories with morphisms *right exact* functors (i.e. functors preserving finite co-limits) and natural isomorphisms. (Recall that a category is essentially small if it is equivalent to a small category.) Note that since abelian categories are finitely cocomplete, there's a full subcategory of **Rex** consisting of small abelian categories, with right exact functors.

Given a **Rex** category $\mathcal{C}$, we may consider its ind-completion $\operatorname{ind}(\mathcal{C}) \in \mathbf{PR}$, which will be **Pr** by construction. Given a **Rex** functor $F : \mathcal{C} \to \mathcal{D}$, we may consider the ind-extension $\operatorname{ind} F : \operatorname{ind}(\mathcal{C}) \to \operatorname{ind}(\mathcal{D})$, which will be co-continuous and compact, i.e., will preserve the subcategory of compact objects. Conversely, given a presentable category $\mathcal{C} \in \mathbf{PR}$, we may consider its subcategory $\operatorname{comp}(\mathcal{C})$ of compact objects, an object of **Rex**. Restricting then to compactly generated categories and compact functors, we find the operations ind and comp define an equivalence

$$\operatorname{ind} : \mathbf{Rex} \overset{\sim}{\longleftrightarrow} \mathbf{Pr} : \operatorname{comp} .$$

It is convenient to go back and forth between the concrete setting of small categories **Rex** and the flexible setting of presentable ones **PR**: the monadic techniques we will develop take **Rex** categories as output, but most naturally produce concrete descriptions of their ind-completions in **PR**. Thanks to the above equivalence, we will move between the settings interchangeably.

Remark 3.1. We will adopt the usual convention of referring to objects in $\operatorname{ind}\mathcal{C}$ for a Rex category $\mathcal{C}$ as ind-objects in $\mathcal{C}$. For instance, an ind-algebra in a monoidal category $\mathcal{C}$ really means an algebra object in $\operatorname{ind}\mathcal{C}$ with respect to the co-continuous extension of the monoidal structure.

We also record here the following proposition, which was explained to us by Daniel Schäppi:

Proposition 3.2. *: If $\mathcal{C}$ is an abelian category and T is a right exact monad on $\mathcal{C}$, then $\mathcal{D} := T - mod_{\mathcal{C}}$ is abelian.*

Proof. The forgetful functor $\mathcal{D} \to \mathcal{C}$ creates all limits that exist in $\mathcal{C}$ (for any monad), in particular finite ones, and all finite colimits (since T preserves finite colimits). Since the forgetful functor is also conservative and $\mathcal{C}$ is abelian, it follows that the comparison morphism

$$coker(ker(f)) \longrightarrow ker(coker(f))$$

is an isomorphism in $\mathcal{D}$, hence $\mathcal{D}$ is abelian. □

3.2. **The Deligne-Kelly tensor product.** For $\mathbf{k}$-linear categories $\mathcal{C}, \mathcal{D}, \mathcal{E}$, let $\operatorname{Bilin}(\mathcal{C} \times \mathcal{D}, \mathcal{E})$ denote the category of $\mathbf{k}$-bilinear functors from $\mathcal{C} \times \mathcal{D}$ to $\mathcal{E}$, preserving finite colimits in each variable separately.

Definition 3.3. The *Deligne-Kelly tensor product* $\mathcal{C} \boxtimes \mathcal{D}$ of $\mathcal{C}, \mathcal{D} \in \mathbf{Rex}$ is uniquely characterized by the natural equivalences,

$$\mathbf{Rex}[\mathcal{C} \boxtimes \mathcal{D}, \mathcal{E}] \simeq \operatorname{Bilin}(\mathcal{C} \times \mathcal{D}, \mathcal{E}) \simeq \mathbf{Rex}[\mathcal{C}, \mathbf{Rex}[\mathcal{D}, \mathcal{E}]].$$

In [Kel89], it is shown that Kelly tensor product equips **Rex** with the structure of a symmetric closed monoidal (2,1)-category, which we denote $\mathbf{Rex}^{\boxtimes}$. The Kelly tensor product extends to **Pr**, where it is characterized by the analogous universal property (with respect to functors preserving colimits in both factors), and the functor ind extends to a symmetric monoidal equivalence $\mathbf{Rex}^{\boxtimes} \simeq \mathbf{Pr}^{\boxtimes}$.

Remark 3.4. It is shown in [LF12] that the Deligne tensor product of abelian categories – when it exists – coincides with the Kelly tensor product, but that the former may not exist in general. This, essentially, is the reason to work with **Rex** and **Pr**, rather than **Ab**.

Proposition 3.5. *The symmetric monoidal ∞-category $\mathbf{Rex}^{\boxtimes} \simeq \mathbf{Pr}^{\boxtimes}$ is closed under small colimits, and the tensor product preserves colimits in each factor separately.*

Proof. In [BKP89] it is shown that **Rex** (equivalently, **Lex**) is the category of 2-modules in **Cat** of a finitary 2-monad T, and as such is closed under arbitrary bicolimits. Because $\mathbf{Rex}^{\boxtimes}$ is *closed* monoidal, the functor $\mathcal{C} \boxtimes -$ has a right adjoint, and therefore commutes with arbitrary colimits. □

In particular it follows that **Rex** $\simeq$ **Pr** satisfies the conditions (*) of [AF12] for the definition of factorization homology (for which a much smaller class of colimits is required, namely the sifted ones).

3.3. **Tensor and braided tensor categories.** It is well known that E_1-, E_2- and framed E_2-algebras, respectively, in $\mathrm{Cat}^\times$ are equivalent to monoidal, braided monoidal and balanced braided monoidal categories. In this section we consider the **k**-linear analogs of these structures, as well as the corresponding notion of rigidity (for a general introduction to rigid tensor categories, see e.g. [EGNO15]).

Definition 3.6. A *tensor category* in **Pr** is an E_1-algebra $\mathcal{A}$ in **Pr**. Similarly a *braided tensor* category in **Pr** is an E_2-algebra in **Pr**.

This definition is a compact formulation of the traditional notions of tensor and braided tensor categories. Since **Pr** is a 2-category, the data of an E_1-algebra consists of a category, a functor of tensor product, and an associator natural isomorphism, which satisfies the so-called "pentagon equation". We will follow the usual convention of dropping explicit mention of associators in formulas, for clarity of exposition.

We will from here forward moreover assume that our tensor categories and braided tensor categories are *rigid*:

Definition 3.7. A tensor category $\mathcal{A}$ is rigid if all *compact* objects of $\mathcal{A}$ are left and right dualizable. A braided tensor category is rigid if its underlying E_1-algebra is rigid.

For later use, we note here that the braiding on a braided tensor category $\mathcal{A}$ endows the iterated tensor functors,

$$T^k : \mathcal{A}^k \to \mathcal{A},$$
$$a_1 \boxtimes \cdots \boxtimes a_k \to a_1 \otimes \cdots \otimes a_k,$$

with the structure of a tensor functor, via the following "shuffle" braiding:

$$J_{\mathbf{a},\mathbf{b}} : a_1 \otimes \cdots \otimes a_k \otimes b_1 \otimes \cdots \otimes b_k \xrightarrow{\simeq} a_1 \otimes b_1 \otimes \cdots \otimes a_k \otimes b_k \tag{3.1}$$

$$J_{\mathbf{a},\mathbf{b}} = \sigma_{a_k,b_{k-1}} \circ \cdots \sigma_{a_3\otimes\cdots\otimes a_k,b_2} \circ \sigma_{a_2\otimes\cdots\otimes a_k,b_1}.$$

Remark 3.8. In order to define a $Disk^2_{or}$-algebra in **Rex**, we require $\mathcal{A}$ to be equipped with a balancing. It is well-known [Sel11] that, having already assumed $\mathcal{A}$ is rigid, this is equivalent to equipping $\mathcal{A}$ with a pivotal structure.

Definition 3.9. A tensor functor $F : \mathcal{A} \to \mathcal{B}$ is called *dominant* if every object of $\mathcal{B}$ appears as a sub-object (equivalently using rigidity, quotient) of an object in the image of F.

Remark 3.10. Note that, by Lemma 2.1 of [BN11], a tensor functor $F : \mathcal{A} \to \mathcal{B}$ is dominant if, and only if, F^R is faithful, i.e. $\mathcal{B}$ should be generated under colimits by the image of $\mathcal{A}$.

Definition 3.11. Let $\mathcal{A}$ be a tensor category in **Pr**.

(1) A (right) $\mathcal{A}$-module category $\mathcal{M}$ for a tensor category $\mathcal{A}$ in **Pr** is a category $\mathcal{M} \in$ **Pr**, together with an action functor,

$$act_{\mathcal{M}} : \mathcal{M} \boxtimes \mathcal{A} \to \mathcal{M},$$
$$m \boxtimes X \mapsto m \otimes X,$$

satisfying standard associativity (pentagon) axioms. A left module category is defined similarly[4].

(2) An $(\mathcal{A},\mathcal{B})$-bimodule category for tensor categories $\mathcal{A}$ and $\mathcal{B}$ is, equivalently, a right module category for $\mathcal{A}^{\otimes-op}\boxtimes\mathcal{B}$ or a left module category for $\mathcal{A}\boxtimes\mathcal{B}^{\otimes-op}$.

(3) For $m\in\mathcal{M}$, we denote by act_m the action functor,

$$\begin{aligned}\mathrm{act}_m : \mathcal{A} &\to \mathcal{M}\\ a &\mapsto m\otimes a.\end{aligned}$$

This functor has a right adjoint,

$$\mathrm{act}_m^R : \mathcal{M}\to\mathcal{A}.$$

(4) We denote by $\underline{\mathrm{End}}_{\mathcal{A}}(m) := \mathrm{act}_m^R(m) = \mathrm{act}_m^R(\mathrm{act}_m(\mathbf{1}))$ the *internal endomorphism algebra of* m, which carries a natural algebra structure.

Proposition 3.12. *Let $\mathcal{A}$ be rigid, $\mathcal{M}$ and $\mathcal{N}$ $\mathcal{A}$-module* **Pr** *categories, and $F:\mathcal{M}\to\mathcal{N}$ an $\mathcal{A}$-module* **Pr** *functor admitting a right (respectively left) adjoint as a plain functor. Then the adjoint admits a canonical $\mathcal{A}$-linear structure.*

Proof. The assertion appears as Corollary 2.13 of [DSPS14] (and as an exercise to the reader in [EO04, 3.3]); see also [BZN09, Lemma 3.5] in the dg setting. The adjoint is automatically (op)lax monoidal, and one shows it is in fact strict by writing objects of $\mathcal{A}$ as filtered colimits of compact, hence dualizable, objects which can be moved across the adjunction. □

Definition 3.13 (See [Ost03, DSPS13])**.** Let $\mathcal{A}$ be an abelian tensor category in **Pr** with abelian module category $\mathcal{M}\in\mathbf{Pr}$. We say that $m\in\mathcal{M}$ is:

(1) An $\mathcal{A}$-*generator*, if act_m^R is faithful.
(2) $\mathcal{A}$-*projective* if act_m^R is colimit-preserving.[5]
(3) An $\mathcal{A}$-*progenerator*, or progenerator for the $\mathcal{A}$-action, if it is an $\mathcal{A}$-projective $\mathcal{A}$-generator.

Proposition 3.14. *Let $\mathcal{A}$ be a rigid tensor category in* **Pr**. *Then the tensor product functor $T:\mathcal{A}\boxtimes\mathcal{A}\to\mathcal{A}$ has a co-continuous right adjoint.*

Proof. Since $\mathcal{A}$ is rigid, the right adjoint T^R is an $\mathcal{A}\boxtimes\mathcal{A}^{\otimes-op}$-module functor. Hence for any $X\in\mathcal{A}$,

$$T^R(X) = T^R((X\boxtimes\mathbf{1}_{\mathcal{A}})\otimes\mathbf{1}_{\mathcal{A}})\cong(X\boxtimes\mathbf{1}_{\mathcal{A}})\otimes T^R(\mathbf{1}_{\mathcal{A}}).$$

Since the tensor product of $\mathcal{A}\boxtimes\mathcal{A}^{\otimes-op}$ is cocontinuous in each variable, T^R is cocontinuous. □

Proposition 3.15. *Let $\mathcal{A}$ be a rigid tensor category, and $\mathcal{M}$ a right $\mathcal{A}$-module. Then the action functor*

$$\mathrm{act}:\mathcal{M}\boxtimes\mathcal{A}\to\mathcal{M}$$

has a cocontinuous right adjoint.

---

[4] Henceforth, "module category" will connote "right module category"

[5] We note that this is equivalent to asking that act_m^R preserves finite colimits, since by construction act_m^R preserves filtered colimits.

Proof. The proposition and its proof carry over without modification from the corresponding assertion in the dg category setting, Proposition D.2.2 of [Gai15]. Namely one can write down explicitly a cocontinuous right adjoint act^R and its adjunction data. The functor act^R is given as a composition

$$\mathcal{M} \xrightarrow{Id_{\mathcal{M}} \boxtimes \mathbf{1}_{\mathcal{A}}} \mathcal{M} \boxtimes \mathcal{A} \xrightarrow{Id_{\mathcal{M}} \boxtimes T^R} \mathcal{M} \boxtimes \mathcal{A} \boxtimes \mathcal{A} \xrightarrow{\mathrm{act} \boxtimes Id_{\mathcal{A}}} \mathcal{M} \boxtimes \mathcal{A}$$

of cocontinuous functors, hence is cocontinuous. □

Definition 3.16. Let $\mathcal{M}, \mathcal{N}$ be right and left module categories for a tensor category $\mathcal{A}$, and let $\mathcal{E} \in \mathbf{Pr}$. Recall that $F : \mathcal{M} \boxtimes \mathcal{N} \to \mathcal{E}$ is called $\mathcal{A}$-*balanced* when equipped with natural isomorphisms,

$$B_{m,X,n} : F(m \otimes X \boxtimes n) \cong F(m \boxtimes X \otimes n),$$

for $m \in \mathcal{M}, X \in \mathcal{A}, n \in \mathcal{N}$, satisfying certain coherences (see [ENO10], Definition 3.1). We denote by $\mathrm{Bal}_{\mathcal{A}}(M \boxtimes N, \mathcal{E})$ the category of balanced functors.

Definition 3.17. The *relative Kelly tensor product* $\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N}$ of left and right module categories $\mathcal{M}$ and $\mathcal{N}$ for a tensor category $\mathcal{A}$ is defined, uniquely[6], by the natural equivalence:

$$\mathbf{Pr}[\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N}, \mathcal{E}] \simeq \mathrm{Bal}_{\mathcal{A}}(\mathcal{M} \boxtimes \mathcal{N}, \mathcal{E}).$$

Existence of relative tensor products in $\mathbf{Pr}^{\boxtimes}$ is an easy consequence of existence of Kelly tensor products, and closure of $\mathbf{Pr}^{\boxtimes}$ under colimits. (See [DSPS14] for an explicit study of balanced tensor products.)

Remark 3.18. One might instead define the relative tensor product as the colimit of the infinite 2-sided bar construction for $\mathcal{M}$ and $\mathcal{N}$. It is not hard to show, along the lines of MacLane's coherence theorem, that the resulting colimit is equivalent to the balanced tensor product defined above: essentially, because $\mathbf{Pr}$ is only a 2-category, the infinite bar construction strictifies after the second step.

Proposition 3.19. *Let $\mathcal{A}$ be a rigid tensor category in* $\mathbf{Pr}$, *and let $\mathcal{M}, \mathcal{N}$ be module categories. Then the right adjoint of the canonical functor*

$$\mu : \mathcal{M} \boxtimes \mathcal{N} \to \mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N}$$

is monadic.

Proof. Recall that $\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N}$ can be realized as the colimit in $\mathbf{PR}$ of the simplicial diagram given by the two-sided bar construction, where the simplices are iterated tensor products $\mathcal{M} \boxtimes \mathcal{A} \cdots \boxtimes \mathcal{A} \boxtimes \mathcal{N}$. Such a colimit of presentable categories with left adjoints can be calculated as the limit in the opposite ∞-category $\mathbf{PR}^R$ of presentable categories with right adjoints over the corresponding cosimplicial diagram of right adjoints. By Propositions 3.14 and 3.15, all functors appearing in the two-sided bar construction have cocontinuous right adjoints, so that we are in fact calculating a limit over a cosimplicial diagram in $\mathbf{PR}$. In particular it follows that μ^R is itself a morphism in $\mathbf{PR}$, i.e., cocontinuous (as well as continuous by virtue of being a right adjoint). The limit in turn is identified with the corresponding limit in the 2-category of categories, i.e., with the category of compatible collections of objects in the cosimplices. However forgetting from compatible collections to objects of the 0-th term is a conservative functor: a map of compatible collections (whose objects are determined by the 0-th term) is an isomorphism if and only if it is so on the 0-th term. Thus μ^R is continuous, cocontinuous and conservative, and thus monadic. $\square$

---

[6] up to a contractible choice of equivalences

Corollary 3.20. *If $\mathcal{A}, \mathcal{M}$ and $\mathcal{N}$ are moreover abelian then so is $\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N}$.*

Proof. Since we are working with compact functors, the monad from the proposition is itself colimit preserving. Thus Proposition 3.2 applies, and when $\mathcal{A}, \mathcal{M}$ and $\mathcal{N}$ are abelian so is $\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N}$. $\square$

4. Barr-Beck reconstruction of module categories

A key tool in our computation is the well-known yoga of monads (= unital algebra objects in categories of endofunctors); in particular, monads arising from adjunctions of tensor functors and module functors. Many of these results are straightforward extensions to **Pr** of the work of Ostrik [Ost03, EO04], in the setting of fusion categories, and [DSPS13] for general finite abelian categories; we are grateful to Ostrik and Snyder for many explanations.

We begin by recalling Beck's monadicity theorem. Let $(L, R) : \mathcal{C} \underset{R}{\overset{L}{\rightleftharpoons}} \mathcal{D}$ be an adjoint pair of functors. The composition $T = R \circ L$ is naturally a monad on $\mathcal{C}$ via the adjunction unit and counit,

$$\eta : \mathrm{id}_{\mathcal{C}} \to R \circ L, \qquad m : R \circ L \circ R \circ L \xrightarrow{\mathrm{id} \otimes \epsilon \otimes \mathrm{id}} R \circ L.$$

Definition 4.1. We denote by T-mod$_{\mathcal{C}}$ *the category of T-modules*[7] *in* $\mathcal{C}$: objects are pairs (X, f) of an object $X \in \mathcal{C}$, and a morphism $f : T(X) \to X$; morphisms from (X, f) to (Y, g) are those $h : X \to Y$ making the obvious diagram commute.

We obtain a functor, $\widetilde{R} : \mathcal{D} \to T - mod_{\mathcal{C}}$, sending $A \in \mathcal{D}$ to $R(A) \in \mathcal{C}$, with its canonical T-action:

$$act : R \circ L \circ R(A) \xrightarrow{\mathrm{id} \otimes \epsilon} R(A).$$

Theorem 4.2 (Barr-Beck theorem)**.** *The functor $\widetilde{R}$ is an equivalence if, and only if:*

- *R is conservative/reflects isomorphism, i.e. $R(X) \cong R(Y) \Rightarrow X \cong Y$.*
- *$\mathcal{D}$ has coequalizers of R-split parallel pairs (those parallel pairs of morphisms in $\mathcal{D}$, which R sends to pairs having a split coequalizer in $\mathcal{C}$), and R preserves those coequalizers.*

Lemma 4.3. *Suppose that $\mathcal{D}$ is abelian. Then R is conservative if, and only if, for any X with $R(X) \cong 0$ we have $X \cong 0$.*

Proof. Suppose we have $f : X \to Y$, and that $R(f)$ an isomorphism. Since R is right exact (in fact cocontinuous) it preserves finite colimits, and since R is a right adjoint, it preserves arbitrary limits; hence $R(\ker f) = \ker R(f) = 0$ and $R(\operatorname{coker} f) = \operatorname{coker} R(f) = 0$, hence $\ker f$ and $\operatorname{coker} f$ are zero, hence f is an isomorphism. $\square$

Remark 4.4. We note that without the abelian assumption, this only gives us that f is monic and epic. Thus to check conservativity of functors in what follows, it will be extremely useful to know the categories involved are abelian – in particular balanced – so that monic and epic maps are isomorphisms.

---

[7] These are sometimes called T-algebras, instead of T-modules.

Theorem 4.5 (Monadicity for module categories)**.** *Let $\mathcal{A}$ be a rigid abelian tensor category in $\mathbf{Pr}$, and let $\mathcal{M} \in \mathbf{Pr}$ be an abelian $\mathcal{A}$-module category with an $\mathcal{A}$-progenerator $m \in \mathcal{M}$. Then we have an equivalence of $\mathcal{A}$-module categories,*

$$\mathcal{M} \simeq \underline{\mathrm{End}}(m)\text{-}\mathrm{mod}_{\mathcal{A}},$$

where $\mathcal{A}$ acts on the right by multiplication.

Proof. We apply Theorem 4.2 to the functor $\mathrm{act}_{\mathcal{M}}^R : \mathcal{A} \to \mathcal{M}$. We need to check that the right adjoint act_m^R is conservative and co-continuous, which is precisely the assumption that m is a pro-generator. Hence act_m^R is monadic. Because $\mathcal{A}$ is rigid, act_m^R carries a canonical module structure, so that the monad $\mathrm{act}_m^R \circ \mathrm{act}_m$ is a module functor, and can therefore be identified with the functor of tensoring with the algebra object $\mathrm{act}_m^R \circ \mathrm{act}_m(\mathbf{1}_{\mathcal{A}}) \cong \underline{\mathrm{End}}(m)$. ☐

Remark 4.6. Note that a *right* $\mathcal{A}$-module category with an $\mathcal{A}$-progenerator is identified with the category of *left*-modules for the internal endomorphism algebra, and vice versa.

There are two senses in which the construction of internal endomorphism algebras is functorial: with respect to $\mathcal{A}$-module functors, and with respect to base change along a dominant tensor functor $F : \mathcal{A} \to \mathcal{B}$.

Theorem 4.7 (Functoriality of monads)**.** *Let $\mathcal{M}$ and $\mathcal{N}$ be $\mathcal{A}$-module categories in $\mathbf{Pr}$, $F : \mathcal{M} \to \mathcal{N}$ an $\mathcal{A}$-module functor, and let $m \in \mathcal{M}$. Then we have a canonical homomorphism of algebras,*

$$\rho_F : \underline{\mathrm{End}}_{\mathcal{A}}(m) \to \underline{\mathrm{End}}_{\mathcal{A}}(F(m)),$$

Proof. We have the following commutative diagram of functors:

$$\begin{tikzcd}
& \mathcal{M} \arrow[dl] \arrow[dd, bend left, "F"] \\
\mathcal{A} \arrow[ur, bend left, "\mathrm{act}_m"] \arrow[dr, bend right, "\mathrm{act}_{F(m)}"'] & \\
& \mathcal{N}. \arrow[ul] \arrow[uu, bend left, "F^R"]
\end{tikzcd}$$

Hence, we define:

$$\rho_F : \underline{\mathrm{End}}(m) = \mathrm{act}_m^R \circ \mathrm{act}_m(\mathbf{1}) \xrightarrow{\eta_F} \mathrm{act}_m^R \circ F^R \circ F \circ \mathrm{act}_m(\mathbf{1})$$

$$\cong \mathrm{act}_{F(m)}^R \circ \mathrm{act}_{F(m)}(\mathbf{1}) = \underline{\mathrm{End}}(F(m)).$$

It is straightforward to see that $\tilde{F}$ is compatible with monadic composition, so defines an algebra homomorphism. ☐

Corollary 4.8. *Suppose that m is an $\mathcal{A}$-progenerator for $\mathcal{M}$, and $F(m)$ is an $\mathcal{A}$-progenerator for $\mathcal{N}$, so that we have equivalences,*

$$\mathcal{M} \simeq \underline{\mathrm{End}}(m)\text{-}\mathrm{mod}_{\mathcal{A}}, \qquad \mathcal{N} \simeq \underline{\mathrm{End}}(F(m))\text{-}\mathrm{mod}_{\mathcal{A}}.$$

Then F is naturally isomorphic to the pull-back functor along ρ_F.

Theorem 4.9 (Monadicity for relative tensor products)**.** *Let $\mathcal{A}$ be a tensor category, and let $\mathcal{M}$, $\mathcal{N}$ be right and left $\mathcal{A}$-module abelian categories, respectively, with $\mathcal{A}$-progenerators $m \in \mathcal{M}$ and $n \in \mathcal{N}$. Then we have equivalences of categories:*

$$\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N} \simeq \underline{\mathrm{End}}(m)\text{-mod}_{\mathcal{N}} \simeq (\underline{\mathrm{End}}(m)\text{-}\underline{\mathrm{End}}(n))\text{-bimod}_{\mathcal{A}}\,.$$

Proof. To begin, we construct a Rex functor, $G : \mathcal{M} \boxtimes \mathcal{N} \to \mathcal{N}$,

$$m' \boxtimes n' \mapsto \underline{\mathrm{Hom}}(m, m') \otimes m'.$$

We note that G carries an evident balancing, $G((m' \otimes a) \boxtimes n') \cong G(m' \boxtimes (a \otimes n'))$:

$$F(\underline{\mathrm{Hom}}(m, m' \otimes a)) \otimes n' \cong F(\underline{\mathrm{Hom}}(m, m') \otimes a) \otimes n' \cong F(\underline{\mathrm{Hom}}(m, m')) \otimes F(a) \otimes n'.$$

This balancing therefore defines an **Pr** functor,

$$G_{\mathcal{A}} : \mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N} \to \mathcal{N}.$$

Moreover $G_{\mathcal{A}}$ carries an action of $\underline{\mathrm{End}}(m)$:

$$\underline{\mathrm{End}}(m) \otimes \underline{\mathrm{Hom}}(m, m') \otimes a \otimes n' \cong \underline{\mathrm{End}}(m) \otimes \underline{\mathrm{Hom}}(m, m') \otimes a \otimes n' \to$$

$$\to F(\underline{\mathrm{Hom}}(m, m') \otimes a) \otimes n'.$$

It can easily be checked that $n' \mapsto m \boxtimes_{\mathcal{A}} n'$ defines a left adjoint $G_{\mathcal{A}}^L$ to $G_{\mathcal{A}}$, and that $G_{\mathcal{A}} G_{\mathcal{A}}^L(n') \cong \underline{\mathrm{End}}(m) \otimes n'$.

Lemma 4.10. *The functor $G_{\mathcal{A}}$ is conservative.*

Proof. We use the abelian property. For this, we note that $G_{\mathcal{A}}$ factors as a composition,

$$\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N} \to Fun_{\mathcal{A}}(\mathcal{M}^{opp}, \mathcal{N}) \xrightarrow{ev_m} \mathcal{N},$$

where the first map is an equivalence using the fact that $\mathcal{M}$ is dualizable as an $\mathcal{A}$-module, and the underlying category of the $\mathcal{A}$-dual is its **Pr** dual – i.e., the **Pr** category with category of compact objects the opposite to the **Rex** category of compact objects in $\mathcal{M}$. (See [Gai15, D.5], [BZN09, 2.9] or [DSPS13, 3.4.9].) The second map is conservative by the assumption that m is an $\mathcal{A}$-generator: an $\mathcal{A}$-functor vanishing on m must vanish identically. Hence $G_{\mathcal{A}}$ is conservative as well. □

Hence to conclude the proof of Theorem 4.9, we may apply Theorem 4.2 to identify $\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N}$ with the category of $\underline{\mathrm{End}}(m)$-modules in $\mathcal{N}$. The second equivalence in the theorem is the observation that, for algebras A and B, an A-B-bimodule is simply a left A-module in the category of right B-modules. □

Corollary 4.11 (Monadicity for base change)**.** *Let $\mathcal{M}$ be an abelian $\mathcal{A}$-module category, $F : \mathcal{A} \to \mathcal{B}$ a dominant tensor functor, and $m \in \mathcal{M}$ a $\mathcal{A}$-progenerator. Then $m \boxtimes_{\mathcal{A}} \mathbf{1}_{\mathcal{B}}$ is a $\mathcal{B}$-progenerator of $\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{B}$, and we have an equivalence of $\mathcal{B}$-module categories,*

$$\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{B} \simeq F(\underline{\mathrm{End}}(m))\text{-mod}_{\mathcal{B}}\,.$$

Proof. This is a special case of Theorem 4.9, simply noting that the functors G and $G_{\mathcal{A}}$ are canonically $\mathcal{B}$-module functors in the case $\mathcal{N} = \mathcal{B}$, regarded as a left $\mathcal{A}$-module through the functor F, and right $\mathcal{B}$-module by the right regular action. □

Remark 4.12. We note that, in the absence of a braiding, the category of $A-B$ bimodules over algebras A and B, internal to a tensor category $\mathcal{A}$, cannot be written as the category of $A \otimes B^{op}$-modules in $\mathcal{A}$, as in the case $\mathcal{A} = \mathrm{Vect}$, simply because there is no natural algebra structure on $A \otimes B^{op}$.

4.1. **Barr-Beck-Tannaka-Krein reconstruction for braided tensor categories.** We shall assume from now on that $\mathcal{A}$ is a rigid braided tensor category. We highlight two important special cases where monadicity applies:

Proposition 4.13. *For any n, the tensor unit $\mathbf{1}_{\mathcal{A}}$ is a progenerator for the n-fold right regular $\mathcal{A}^{\boxtimes n}$-action.*

Proof. This reduces immediately to Proposition 3.14 □

Remark 4.14. It is possible to describe $\underline{\mathrm{End}}_{\mathcal{A}^{\boxtimes 2}}(\mathbf{1}_{\mathcal{A}})$ explicitly, via the co-end construction; we have:

$$\underline{\mathrm{End}}({}_{\mathcal{A}}\mathbf{1}_{\mathcal{A}}) = \left(\bigoplus_{V \in \mathrm{comp}\,\mathcal{A}} V^* \boxtimes V\right) \Big/ \langle \mathrm{Im}(\mathrm{id}_{W^*} \boxtimes \phi - \phi^* \boxtimes \mathrm{id}_V) \mid \phi : V \to W \rangle. \tag{4.1}$$

If $\mathcal{A}$ is semi-simple, we may make this more explicit by choosing a representative X of every simple isomorphism class of object and write simply,

$$\underline{\mathrm{End}}_{\mathcal{A}^{\boxtimes 2}}(\mathbf{1}_{\mathcal{A}}) \cong \bigoplus_X X^* \boxtimes X.$$

When expressed through the canonical maps $\iota_U : U^* \boxtimes U \to \underline{\mathrm{End}}({}_{\mathcal{A}}\mathbf{1}_{\mathcal{A}})$, the algebra structure is the tautological one, composed with the braiding:

$$V^* \boxtimes V \otimes W^* \boxtimes W = (V^* \otimes W^*) \boxtimes (V \otimes W) \xrightarrow{\sigma_{V^*,W^*}} (W^* \otimes V^*) \boxtimes (V \otimes W) \xrightarrow{\iota_{V \otimes W}} \underline{\mathrm{End}}({}_{\mathcal{A}}\mathbf{1}_{\mathcal{A}}).$$

Definition 4.15. Let $\mathcal{O}(\mathcal{A}) := T(\underline{\mathrm{End}}_{\mathcal{A}^{\boxtimes 2}}(\mathbf{1}_{\mathcal{A}}))$ denote the algebra in $\mathcal{A}$ obtained by applying the tensor product $T : \mathcal{A} \boxtimes \mathcal{A} \to \mathcal{A}$.

As a special case of monadicity for base change, we have:

Theorem 4.16 (Monadicity for traces)**.** *We have an equivalence of categories,*

$$\mathcal{A} \underset{\mathcal{A} \boxtimes \mathcal{A}}{\boxtimes} \mathcal{A} \simeq \mathcal{O}(\mathcal{A})\text{-}\mathrm{mod}_{\mathcal{A}}.$$

Clearly, the description of $\underline{\mathrm{End}}_{\mathcal{A}^{\boxtimes 2}}(\mathbf{1}_{\mathcal{A}})$ implies a similar presentation,

$$\mathcal{O}(\mathcal{A}) = \left(\bigoplus_{V \in \mathcal{A}} V^* \otimes V\right) \Big/ \langle \mathrm{Im}(\mathrm{id}_{W^*} \otimes \phi - \phi^* \otimes \mathrm{id}_V) \mid \phi : V \to W \rangle, \tag{4.2}$$

and in the case $\mathcal{A}$ is semi-simple, we may write:

$$\mathcal{O}(\mathcal{A}) \cong \bigoplus_X X^* \otimes X.$$

The multiplication map $\mu : \mathcal{O}(\mathcal{A}) \otimes \mathcal{O}(\mathcal{A}) \to \mathcal{O}(\mathcal{A})$ may be expressed entirely internally to $\mathcal{A}$. Using the canonical maps $\iota_V : V^* \otimes V \to \mathcal{O}(\mathcal{A})$, we may write the multiplication as follows:

$$(V^* \otimes V) \otimes (W^* \otimes W) \xrightarrow{\sigma_{V^* \otimes V, W^*}} (W^* \otimes V^*) \otimes (V \otimes W) \xrightarrow{\iota_{V \otimes W}} \mathcal{O}(\mathcal{A}).$$

Remark 4.17. When $\mathcal{A} = \operatorname{Rep_q} G$, the algebra $\mathcal{O}(\mathcal{A})$ goes by many names: it has appeared in the literature as the "equivariantized quantum coordinate algebra" of Majid, "the reflection equation algebra" (usually, in the case of GL_N), and the "quantum loop algebra" of Alekseev and Schomerus, and finally is isomorphic via the Rosso form to the locally finite subalgebra of $U_q\mathfrak{g}$.

5. Computing factorization homology of punctured surfaces

This section contains the first main result of this paper, that is an explicit computation of the factorization homology of a braided tensor category on any punctured surface.

5.1. **The distinguished object.** Recall from Section 2.2 that the inclusion

$$\emptyset \longrightarrow M$$

of the empty manifold into any surface induces a canonical functor

$$\mathrm{Vect}_{\mathbf{k}} \longrightarrow \int_M \mathcal{A}.$$

Definition 5.1. The image of the tensor unit $\mathbf{k} \in \mathrm{Vect}$ under the above functor will be called the distinguished object or quantum structure sheaf, and denoted by $\mathcal{O}_{\mathcal{A},M} \in \int_M \mathcal{A}$.

We note that by construction, the distinguished object is mapped to the distinguished object by the functor on factorization homology induced by an embedding of manifolds. In this section we explicitly compute the functor associated to certain particular embeddings:

Proposition 5.2. *Let $X = (\mathbb{R}^2)^{\sqcup k}$ be a disjoint union of k 2-disks. We have:*

1. *We have an isomorphism $\mathcal{O}_{\mathcal{A},X} \cong \mathbf{1}_{\mathcal{A}^{\boxtimes k}}$.*
2. *The functor on factorization homology induced by any embedding $X \to \mathbb{R}^2$ is isomorphic to the tensor functor $\mathcal{A}^{\boxtimes k} \to \mathcal{A}$.*
3. *Any two embeddings of X into a path connected manifold M give rise to isomorphic functors, and any such factors through the tensor functor $\mathcal{A}^{\boxtimes k} \to \mathcal{A}$.*

Proof. Parts (1) and (2) are contained in the equivalence between locally constant factorization algebras on $\mathbb{R}^2$ and E_2-algebras. Part (3) follows from the fact that $\mathrm{Emb}(X, M)$ is path connected, and the fact that any such embedding can be factored through an embedding of X into a bigger disk. ☐

Proposition 5.3. *Let $M = X \underset{P\times\mathbb{R}}{\sqcup} Y$ be a collar gluing of two manifolds X and Y, along a codimension 1 sub-manifold P with a choice of a trivialization $N \cong P \times \mathbb{R}$ of an open tubular neighborhood N of P.*

1. *The map $X \sqcup Y \to M$ given by the composition $X \sqcup Y \cong M \backslash N \hookrightarrow M$ induces the canonical functor from absolute to relative tensor product:*
$$\int_X \mathcal{A} \boxtimes \int_Y \mathcal{A} \longrightarrow \int_X \mathcal{A} \underset{\int_N \mathcal{A}}{\boxtimes} \int_Y \mathcal{A}.$$
2. *Moreover, we have a canonical isomorphism:*
$$\mathcal{O}_{\mathcal{A},M} \cong \mathcal{O}_{\mathcal{A},X} \underset{\int_N \mathcal{A}}{\boxtimes} \mathcal{O}_{\mathcal{A},Y}.$$

Proof. Part (1) can be extracted form the proof of the excision property. Since $\emptyset$ is initial, the maps

$$\emptyset \to M$$

and

$$\emptyset \cong \emptyset \sqcup \emptyset \to X \sqcup Y \to M$$

coincide, which together with (1) implies (2). □

5.2. **Moduli algebras.** Let S be a punctured surface. We define the following algebra:

Definition 5.4. The moduli algebra of S is

$$A_S := \underline{\mathrm{End}}_{\mathcal{A}}(\mathcal{O}_{\mathcal{A},S}).$$

The main goal of this section is to describe a combinatorial and explicit presentation of A_S, and to use it to identify $\int_S \mathcal{A}$ with the category of A_S-modules in $\mathcal{A}$.

Punctured surfaces may be indexed by combinatorial data called *gluing patterns* P. In this section, we will give explicit presentations of algebras a_p in $\mathcal{A}$, whose categories $a_p\text{-mod}_{\mathcal{A}}$ of modules in $\mathcal{A}$ describe the factorization homology of the associated marked, punctured surface $\Sigma(P)$.

Definition 5.5. A *gluing pattern* is a bijection,

$$P : \{1, 1', \ldots, n, n'\} \xrightarrow{\sim} \{1, \ldots 2n\},$$

such that $P(i) < P(i')$, for all $i = 1, \ldots, n$. We highlight for future use the following gluing patterns:

$$Ann(1,1') := (1,2), \qquad T^2 \backslash D^2(1,1',2,2') := (1,3,2,4),$$

and by convention, we allow the null gluing pattern $D^2 : \emptyset \to \emptyset$.

Remark 5.6. The set of such gluing patterns, modulo cyclic permutations, is clearly in bijection with the set of ribbon graphs with a single vertex; the indexing we use is most convenient for writing down explicit relations.



FIGURE 1. The punctured surface $\Sigma(P)$ constructed from a gluing pattern P.

A gluing pattern P determines a surface $\Sigma(P)$, with a single marked boundary interval 0, as follows: we begin with a disk D^2 with $2n+1$ boundary intervals labeled $0, 1, \ldots 2n$, and then glue each of n handles H_i, with marked intervals, i and i' to $P(i)$ and $P(i')$, respectively.

Definition 5.7. We say that handles H_i and H_j, with $i < j$ are:

- *positively linked* if $P(i) < P(j) < P(i') < P(j')$,
  negatively linked if $P(j) < P(i) < P(j') < P(i')$,
- *positively nested* if $P(i) < P(j) < P(j') < P(i')$,
  negatively nested if $P(j) < P(i) < P(i') < P(j')$,
- *positively unlinked* if $P(i) < P(i') < P(j) < P(j')$,
  negatively unlinked if $P(j) < P(j') < P(i) < P(i')$,

where the sign is $(+)$ if $i < j$, and $(-)$ if $i > j$.

Definition 5.8. We define the crossing morphisms,

$$L_{ij}, N_{ij}, U_{ij} : \mathcal{O}(\mathcal{A})^{(i)} \otimes \mathcal{O}(\mathcal{A})^{(j)} \to \mathcal{O}(\mathcal{A})^{(j)} \otimes \mathcal{O}(\mathcal{A})^{(i)},$$

as follows:


$\mathcal{O}(\mathcal{A})^{(j)}$
$\mathcal{O}(\mathcal{A})^{(i)}$
$\mathcal{O}(\mathcal{A})^{(i)}$
$\mathcal{O}(\mathcal{A})^{(j)}$

Linked crossing L_{ij}


$\mathcal{O}(\mathcal{A})^{(j)}$
$\mathcal{O}(\mathcal{A})^{(i)}$
$\mathcal{O}(\mathcal{A})^{(i)}$
$\mathcal{O}(\mathcal{A})^{(j)}$

Nested crossing N_{ij}


$\mathcal{O}(\mathcal{A})^{(j)}$
$\mathcal{O}(\mathcal{A})^{(i)}$
$\mathcal{O}(\mathcal{A})^{(i)}$
$\mathcal{O}(\mathcal{A})^{(j)}$

Unlinked crossing U_{ij}

Remark 5.9. The unlinked crossing operator U_{ij} is nothing but the braiding of $\mathcal{A}$ applied to $\mathcal{O}(\mathcal{A})^{(i)} \otimes \mathcal{O}(\mathcal{A})^{(j)}$.

Definition 5.10. For a gluing pattern of rank n, we define the algebra a_P to be the object $\mathcal{O}(\mathcal{A})^{\otimes n}$ in $\mathcal{A}$, with multiplication on each pair of factors $\mathcal{O}^{(i)} \otimes \mathcal{O}^{(j)}$, $i < j$:

$$\mathcal{O}^{(i)} \otimes \mathcal{O}^{(j)} \otimes \mathcal{O}^{(i)} \otimes \mathcal{O}^{(j)} \xrightarrow{\mathrm{id} \otimes C_{ij} \otimes \mathrm{id}} \mathcal{O}^{(i)} \otimes \mathcal{O}^{(i)} \otimes \mathcal{O}^{(j)} \otimes \mathcal{O}^{(j)} \xrightarrow{m \otimes m} \mathcal{O}^{(i)} \otimes \mathcal{O}^{(j)},$$

where $C_{ij} = L_{ij}^{\pm 1}$ (resp. $N_{ij}^{\pm 1}, U_{ij}^{\pm 1}$), if H_i and H_j are $(\pm)$-linked (resp. $(\pm)$-nested, $(\pm)$-unlinked), and m denotes the multiplication on $\mathcal{O}$.

The following is the main result of this section:

Theorem 5.11. *We have an isomorphism of algebras $A_{\Sigma(P)} \cong a_P$, and an equivalence of categories,*

$$\int_{\Sigma(P)} \mathcal{A} \simeq a_P\text{-mod}_{\mathcal{A}} \simeq A_{\Sigma(P)}\text{-mod}_{\mathcal{A}}$$

Proof. We may deform the attaching disk of $\Sigma(P)$ in such a way that the 0-marked interval is on the right, and all other marked intervals are on the left.

We have $\int_D \mathcal{A} \simeq \mathcal{A}$ as a category; however the markings on D induce a $\mathcal{A}^{\boxtimes 2n}$-$\mathcal{A}$-bimodule structure,

$$(a_1 \boxtimes \cdots \boxtimes a_{2n}) \boxtimes b \boxtimes c \mapsto a_1 \otimes \cdots \otimes a_{2n} \otimes b \otimes c,$$

which bimodule we denote by ${}_{2n}\mathcal{A}_{\mathcal{A}}$.

For each handle H_i, we have $\int_{H_i} \mathcal{A} \simeq \mathcal{A}_{\mathcal{A}\mathcal{A}}$, the category $\mathcal{A}$ with its right regular $\mathcal{A} \boxtimes \mathcal{A}$-module structure.

Likewise, given n handles $H_1, \dots H_n$ we have $\int_{H_1 \cup \dots \cup H_n} \mathcal{A} \simeq \mathcal{A}^{\boxtimes n}$ as a category. We make this a right $\mathcal{A}^{\boxtimes 2n}$-module using P:

$$(a_1 \boxtimes \dots \boxtimes a_{n'}) \boxtimes (b_1 \boxtimes \dots \boxtimes b_{2n}) \mapsto (a_1 \otimes b_{P(1)}) \boxtimes \dots \boxtimes (a_{n'} \otimes b_{P(n')}).$$

We denote this right $\mathcal{A}^{\boxtimes 2n}$-module by $\mathcal{A}^P$.

We note that the module structures on A^P and ${}_{2n}\mathcal{A}_{\mathcal{A}}$ are precisely those induced by the markings on the left hand side of Figure 1. Thus, by the excision property for factorization homology, we have:

$$\int_{\Sigma(P)} \mathcal{A} \simeq \mathcal{A}^P \underset{\mathcal{A}^{\boxtimes 2n}}{\boxtimes} {}_{2n}\mathcal{A}_{\mathcal{A}}. \tag{5.1}$$

Let $\tau_P \in S_{2n}$ be the permutation obtained by precomposing P with the map

$$\{1, \dots, 2n\} \to \{1, 1', \dots, n, n'\}$$

defined by

$$i \mapsto \begin{cases} i/2 & \text{if i is even} \\ (i/2)' & \text{if i is odd} \end{cases}$$

Applying Proposition 4.13, we may identify $\mathcal{A}^P$ with the category of modules in $\mathcal{A}^{\boxtimes 2n}$ for an algebra $\underline{\mathrm{End}}(\mathbf{1}_{\mathcal{A}\mathcal{A}})^P$ obtained by applying τ_P to $\underline{\mathrm{End}}(\mathbf{1}_{\mathcal{A}\mathcal{A}})^{\boxtimes n}$.

The bimodule ${}_{2n}\mathcal{A}_{\mathcal{A}}$ is simply that induced by the iterated tensor product functor $T^{2n} : \mathcal{A}^{\boxtimes 2n} \to \mathcal{A}$, which itself carries the structure of a tensor functor using formula (3.1). Hence we may apply Theorem 4.11 to conclude,

$$\int_{\Sigma(P)} \mathcal{A} \simeq T^{2n}(\underline{\mathrm{End}}(\mathbf{1}_{\mathcal{A}\mathcal{A}})^P)\text{-mod}_{\mathcal{A}}, \tag{5.2}$$

as a right $\mathcal{A}$-module category.

Note that equations (5.1) and (5.2), combined with Theorem 4.11, yield an isomorphism,

$$\widetilde{a_P} \cong A_S.$$

Let us now explain how to identify $\tilde{a}_P := T^{2n}(\underline{\mathrm{End}}(\mathbf{1}_{\mathcal{A}\mathcal{A}})^P)$ with the algebra a_P. First, we consider the subalgebras $\mathcal{O}(\mathcal{A})^{(i,i')} = \underline{\mathrm{End}}(\mathbf{1}_{\mathcal{A}_{P(i)}\mathcal{A}_{P(i')}})$, and their images $\mathcal{O}(\mathcal{A})^{(i)} := T(\mathcal{O}(\mathcal{A})^{(i,i')})$. The multiplication induces an isomorphism,

$$m : \mathcal{O}(\mathcal{A})^{(1)} \otimes \dots \otimes \mathcal{O}(\mathcal{A})^{(n)} \to \tilde{a}_p,$$

on the level of objects. It remains only to compute the pairwise cross relations between factors. Note that $\mathcal{O}^{(i,i')}$ and $\mathcal{O}^{(j,j')}$ commute in $\mathcal{A}^{\boxtimes 4}$, because they occupy different tensor factors. Hence we have the following commutative diagram:

(5.3)
$$\begin{tikzcd}
 & T^4(\mathcal{O}(\mathcal{A})^{(i,i')}\otimes\mathcal{O}(\mathcal{A})^{(j,j')}) \arrow[ddr, "T^4(m)"] & \underset{can}{=} & T^4(\mathcal{O}(\mathcal{A})^{(j,j')}\otimes\mathcal{O}(\mathcal{A})^{(i,i')}) \arrow[ddl, "T^4(m)"'] & \\
\mathcal{O}(\mathcal{A})^{(i)}\otimes\mathcal{O}(\mathcal{A})^{(j)} \arrow[ur, "J_{ij}"] \arrow[drr, "m"'] & & & & \mathcal{O}(\mathcal{A})^{(j)}\otimes\mathcal{O}(\mathcal{A})^{(i)} \arrow[ul, "J_{ji}"'] \arrow[dll, "m"] \\
 & & a_P & &
\end{tikzcd}$$

The ordering on the tensor factors in $\mathcal{A}^P$, and hence the value of J_{ij}, depends on the gluing pattern. For example, in the positively linked case, as indicated in Figure 2, we have $J_{12} = \mathrm{id}\otimes\sigma\otimes\mathrm{id}$, and $J_{21} = (\sigma\otimes\sigma)\circ(\mathrm{id}\otimes\sigma\otimes\mathrm{id})$, hence:

$$m|_{\mathcal{O}^{(2)}\otimes\mathcal{O}^{(1)}} = m|_{\mathcal{O}^{(1)}\otimes\mathcal{O}^{(2)}}\circ J_{12}^{-1}J_{21} = m|_{\mathcal{O}^{(1)}\otimes\mathcal{O}^{(2)}}\circ L_{12}^{+},$$

as claimed. The other five cases follow a similar computation.



FIGURE 2. In the linked case, we have $P(1,1',2,2') = (1,3,2,4)$.

$\square$

Definition 5.12. The *disjoint union* of gluing patterns $P : \{1,1',\ldots,n,n'\}\to\{1,\ldots 2n\}$ and $Q : \{1,1',\ldots,m,m'\}\to\{1,\ldots,2m\}$ is

$$P\sqcup Q : \{1,1',\ldots,m+n,(m+n)'\}\to\{1,\ldots,2(m+n)\},$$

$$i,i'\mapsto\begin{cases} P(i),P(i'), & \text{if } i\le n\\ Q(i-n)+2n, Q((i-n)')+2n, & \text{if } i>n\end{cases}$$

We record the following useful corollary.

Corollary 5.13. *Let $P = P_1 \amalg P_2$ be the disjoint union of two gluing pattern. The algebra a_P is the braided tensor product of a_{P_1} and a_{P_2}.*

Proof. By construction, any pair of a handle in P_1 and a handle in P_2 is unlinked, hence the cross relations between the corresponding $\mathcal{O}(\mathcal{A})$ factors are those of the braided tensor product. The result thus follows from the hexagon axioms. $\square$

5.3. **Examples.**

5.3.1. *Reflection equation algebra from the annulus.* The braided dual, or reflection equation algebra, of a quasi-triangular Hopf algebra H is a certain modification of the Hopf dual of H, which is equivariant with respect to the H-adjoint action and is canonically identified with $\mathcal{O}_{H\text{-mod}}$. Donin–Kulish–Mudrov proved in [DKM03] that the reflection equation algebra universally provides representations of the braid group of the annulus, extending the representations of the planar braid group attached to H.

The annulus is homeomorphic to $\Sigma(Ann)$, where we recall that $Ann(1) = 1, Ann(1') = 2$. Thus, we have $a_{Ann} \cong \mathcal{O}(\mathcal{A})$, and hence:



Figure 3. The annulus constructed from a gluing pattern yields $a_{Ann} \cong \mathcal{O}(\mathcal{A})$. The pair of pants yields the braided tensor product $a_{Pan} \cong \mathcal{O}(\mathcal{A})\widetilde{\otimes}\mathcal{O}(\mathcal{A})$.

Theorem 5.14. *There is an equivalence of categories between factorization homology of the annulus with coefficients in H-mod, and the category of H-equivariant modules over the reflection equation algebra of H.*

Choosing bases for objects V, W, the commutation relations between generators $(a_V)^i_j := v^i \otimes v_j$ and $(a_W)^k_l = w^k \otimes w_l$ of $\mathcal{O}(\mathcal{A})$ may be written as follows:

$$\sum_{k,l,m,p} R^{ij}_{kl}(a_V)^l_m R^{mk}_{op}(a_W)^p_r = \sum_{s,t,u,v} (a_W)^i_s R^{sj}_{tu}(a_V)^u_v R^{vt}_{or},$$

or in matrix notation,

$$R_{21}A_V R_{12}A_W = A_W R_{21}A_V R_{12}. \tag{5.4}$$

Remark 5.15. Equation (5.4) is known as the reflection equation, and hence the name reflection equation algebra. In the case $H = U_q(\mathfrak{g})$, the reflection equation algebra is the algebra $\mathcal{O}_q(G)$ which quantizes the Semenov-Tian-Shansky Poisson bracket on G (and should not be confused with the FRT algebra, the Hopf dual to $U_q(\mathfrak{g})$, often denote by the same symbol, of which it is a co-cycle twist [VV10].

5.3.2. *Iterated braided tensor products from pairs of pants.* In [DKM03], Donin–Kulish–Mudrov observed that the nth braided tensor power of the reflection equation algebra provides representations of the braid group of an n-punctured disk. This corresponds to the gluing pattern made of one disk with n unlinked handles. For $n = 2$ this is a pair of pants which is depicted on Figure 3.

Theorem 5.16. *There is an equivalence of categories between factorization homology of n-punctured disk with coefficients in H-mod, and the category of H-equivariant modules over the nth braided tensor power of the reflection equation algebra of H.*

5.3.3. *The elliptic double from the punctured torus.* The punctured torus $T^2\backslash D$ is homeomorphic to $\Sigma(P)$, where $P(1) = 1, P(1') = 3, P(2) = 2, P(2') = 4$.



Figure 4. The punctured torus constructed from a gluing pattern yields $A_{T^2\backslash D} \cong \mathcal{D}_{\mathcal{A}}$.

Hence $A_{T^2\backslash D} \cong \mathcal{D}_{\mathcal{A}}$, where $\mathcal{D}_{\mathcal{A}}$ is the algebra in $\mathcal{A}$ which as an object is $\mathcal{O}(\mathcal{A}) \otimes \mathcal{O}(\mathcal{A})$, and the cross relations of Theorem 5.11 coincide with those of the so-called elliptic double introduced in [BJ14], which provides representations of the braid group of the punctured torus. It is proven there that in the case H is factorizable, $\mathcal{D}_{\mathcal{A}}$ is isomorphic to the so-called Heisenberg double of H. In particular, if $H = U_q\mathfrak{g}$, this is isomorphic in turn to the algebra $\mathcal{D}_q(G)$ of quantum differential operators on G. This algebra has been studied in various context: it is a simultaneous flat deformation of $\mathcal{D}(\mathfrak{g})$ and $\mathcal{D}(G)$ and a quantization of a certain Poisson structure on $G \times G$ thought of as a Poisson-Lie analog of the cotangent bundle of G [STS94], which is one of the building block of the Poisson structure on character varieties (see Section 6).

Theorem 5.17. *There is an equivalence of categories between factorization homology of the punctured torus with coefficients in $\mathrm{Rep}_\mathrm{q}\, G$, and the category of $\mathcal{D}_q(G)$-modules in $\mathrm{Rep}_\mathrm{q}\, G$.*

In order to write this more explicitly, we may choose bases for V and W. Let A_V, A_W denote the $N \times N$ (resp. $M \times M$) matrix $(A_V)^k_l = v^k \boxtimes v_l$, (resp. $(A_W)^m_n = w^m \boxtimes w_n$), with the entries regarded as elements of $\mathcal{O}_{13}$. Let D_V and D_W denote the same matrices, but with elements regarded instead in $\mathcal{O}_{24}$. We can write the commutation relations in matrix form:

$$A_V R_{V,W} D_W = R_{V,W} D_W R_{W,V} A_V R_{V,W},$$

or even more explicitly,

$$\sum_{j,m} a^i_j R^{jk}_{lm} \partial^m_n = \sum_{o,p,r,t,u,v} R^{ik}_{op} \partial^p_r R^{ro}_{tu} a^u_v R^{vt}_{ln},$$

where we have omitted the labels V, W on R-matrices for ease of notation.

5.3.4. *The r-punctured surface $\Sigma_{g,r}$ of genus g.* The g-fold disjoint union of the once-punctured torus gluing pattern yields a gluing pattern for the once-punctured surface of genus g. The disjoint union of this with the $r-1$-fold disjoint union of the annular gluing pattern yields a gluing pattern for $\Sigma_{g,r}$, the r-punctured genus g surface. Accordingly, we have

$$\int_{\Sigma_{g,r}} \mathcal{A} \simeq \mathcal{D}_{\mathcal{A}}^{\otimes g} \otimes \mathcal{O}(\mathcal{A})^{\otimes r-1}\text{-mod}_{\mathcal{A}},$$

where $\otimes$ denotes the braided tensor product in $\mathcal{A}$.



FIGURE 5. The r-punctured, genus g surface from a gluing pattern yields the iterated tensor product of g copies of $\mathcal{D}_{\mathcal{A}}$ and $r-1$ copies of $\mathcal{O}(\mathcal{A})$.

5.4. **Mapping class group actions.** Let $f, g : M \to N$ be diffeomorphisms of manifolds M and N, and suppose that $\eta : f \to g$ is an isotopy. Then η yields an isomorphism of functors, $\eta_* : f_* \xrightarrow{\simeq} g_*$, intertwining the canonical isomorphisms $\iota_f : f_*(\mathcal{O}_{\mathcal{A},M}) \to \mathcal{O}_{\mathcal{A},N}$ and $\iota_g : g_*(\mathcal{O}_{\mathcal{A},M}) \to \mathcal{O}_{\mathcal{A},N}$.

Suppose further that M and N have marked boundary intervals I_M and I_N, respectively, that f and g are maps of pairs, so that $f(I_M) = I_N$ and $g(I_M) = I_N$, and finally that η is an isotopy of maps of pairs, i.e. $\eta_t(I_M) = I_N$ for all t.

Under these assumptions, $\int_M \mathcal{A}$ and $\int_N \mathcal{A}$ are endowed with $\mathcal{A}$-module structures, we have $\mathcal{A}$-module equivalences,

$$f_*, g_* : \int_M \mathcal{A} \to \int_N \mathcal{A}.$$

Along the lines of Proposition 4.7, we have canonical isomorphisms of algebras,

$$\widetilde{f_*}, \widetilde{g_*} : A_M = \underline{\mathrm{End}}_{\mathcal{A}}(\mathcal{O}_{\mathcal{A},M}) \to A_N = \underline{\mathrm{End}}_{\mathcal{A}}(\mathcal{O}_{\mathcal{A},N}),$$

induced by evident isomorphisms of functors,

$$\theta_f, \theta_g : tr_m \xrightarrow{\simeq} tr_n,$$

induced by ι_f, ι_g respectively. Finally, since η_* is canonically an isomorphism of $\mathcal{A}$-module functors, intertwining ι_f and ι_g, we have that θ_f, θ_g, and hence, $\widetilde{f_*}, \widetilde{g_*} : A_M \to A_N$ coincide.

Hence, taking $M = N$, we obtain an action of the mapping class group of M on the algebra A_M.

Remark 5.18. The mapping class group is the truncation $\pi_0 \operatorname{Diff}(M, M)$ of the ∞-groupoid of diffeomorphisms of M, isotopies, isotopies of isotopies, and so on. Clearly, the 2-group $\pi_{\leq 1} \operatorname{Diff}(M, M)$ acts on the category $\int_M \mathcal{A}$ itself. However the simple nature of diffeomorphism groups of surfaces allows one to reduce this action to that of the mapping class group, endowed in the case of genus 0 and 1 with some additional commuting operators (an action of homotopy type of $SO(3)$ and of $S^1 \times S^1$, respectively).

In the case $\mathcal{A} = \operatorname{Rep_q} G$, one recovers this way the mapping class group action on moduli algebras [AS96] and the quantum Fourier transform on the algebra $\mathcal{D}_q(G)$ of differential operators [BJ14]. Indeed, it is proved in *loc. cit.* that there is an action of the universal central extension $\widetilde{SL_2(\mathbb{Z})}$ of the modular group on $\mathcal{D}_q(G)$ by algebra automorphism. Recall that $\widetilde{SL_2(\mathbb{Z})}$ is generated by X, Y with relations

$$X^4 = (XY)^3 \qquad\qquad (X^2, Y) = 1$$

Using the presentation given in Section 5.3.3, the action of the standard generators X, Y is given by:

$$\begin{aligned} X \cdot A_V &= D_V & \qquad X \cdot D_V &= D_V A_V^{-1} D_V^{-1} \\ Y \cdot A_V &= A_V & \qquad Y \cdot D_V &= D_V A_V^{-1}. \end{aligned}$$

We call the automorphism induced by the action of X the "quantum Fourier transform" because it degenerates to the classical Fourier transform on $D(\mathfrak{g})$.

5.5. **Braid group representations.** Fix an embedding $\iota : (\mathbb{R}^2)^{\sqcup k} \hookrightarrow M$. It follows from the formalism of factorization homology that we have an action of the fundamental group $\pi_1(\operatorname{Emb}((\mathbb{R}^2)^{\sqcup k}, M), \iota)$ on the functor

$$F_\iota : \mathcal{A}^{\boxtimes k} \to \int_M \mathcal{A},$$

which is clearly S_k-equivariant. Together with the fact that F_ι factor through the multiplication of $\mathcal{A}$, for each γ in the braid group $B_k(M)$ of M we obtain natural isomorphisms,

$$F_\iota(V_1 \otimes \cdots \otimes V_k) \xrightarrow{\cong} F_\iota(V_{\bar{\gamma}(1)} \otimes \cdots \otimes V_{\bar{\gamma}(k)}),$$

where $\bar{\gamma}$ is the image of γ through $B_k(M) \to S_k$.

This recovers the surface braid group representations obtained in [BJ14, DKM03, Jor09] and generalizes those to higher genus surfaces.

6. CHARACTER VARIETIES AND QUANTIZATION

In this section we explain how our formalism recovers the classical character stack and the character variety, and provides a quantization of the Atiyah–Bott Poisson structure on the latter.

6.1. **Classical character varieties from** Rep G**.** Recall that $\underline{\mathrm{Ch}}_G(S)$ and $\mathrm{Ch}_G(S)$ are respectively the character stack and the character variety defined in the introduction. If S is a connected surface of genus g with $n > 0$ points removed, then $\pi_1(S)$ is the free group on $2g + n - 1$ generators, hence

$$\underline{\mathrm{Ch}}_G(S) \cong [G^{2g+n-1}/G],$$

where the quotient is taken with respect to the diagonal adjoint action. In that case, the category of quasi-coherent sheaves on $\underline{\mathrm{Ch}}_G(S)$ is easily identified with the category of $\mathcal{O}(G^{2g+n-1})$-modules in Rep G. Comparing with Theorem 5.11, we have:

Theorem 6.1. *If S is a punctured surface, then we have an equivalence of categories,*

$$\mathrm{QCoh}(\underline{\mathrm{Ch}}_G(S)) \simeq \int_S \mathrm{Rep}\ G,$$

between the category of quasi-coherent sheaves on $\underline{\mathrm{Ch}}_G(S)$, and the factorization homology of S with coefficients in Rep G*.*

There is a natural identification $\underline{\mathrm{Ch}}_G(S) \cong Map(S, BG)$ where BG is the classifying stack of G. We denote by $\mathrm{Rep}_{\hbar}\,G$ the formal version of $\mathrm{Rep_q}\,G$, defined as in Section 1.3.2 but using the formal quantum group $U_\hbar(\mathfrak{g})$ instead. Hence, if X is a topological space of dimension $0, 1, 2$, factorization homology with coefficients in $\mathrm{Rep}_{\hbar}\,G$ should be thought of as a quantization of the functor

$$QCoh(Map(-, BG)).$$

In [PTVV11] (see also [Toë14]) the authors prove that the classifying stack BG has a canonical 2-shifted symplectic structure induced by the Killing form on $\mathfrak{g}$. It gives rise by integration/pull-back to a 0-shifted symplectic structure on $\underline{\mathrm{Ch}}_G(S)$ which is a stacky version of the Poisson structure on the affine algebraic variety $\mathrm{Ch}_G(S)$. The category $\int_S \mathcal{A}$ is a quantization of this structure.

The main goal of this section is to show directly that our constructions also provide a quantization of the Poisson algebra of functions on the categorical quotient $\mathrm{Ch}_G(S)$, in the sense of deformation quantization.

6.2. **Gluing patterns, fat graphs and Poisson structures.** Let S be a punctured surface and choose a gluing pattern P for S with n handles. The category $\mathrm{Rep}_{\hbar}\,G$ has a tautological strict tensor functor to the category of vector spaces, hence the algebra a_P can be identified with an associative algebra, under this functor. A gluing pattern can be viewed as an instance of a ciliated fat graph in the sense of [FR99], with only one vertex. Recall that G has a standard Poisson Lie group structure. To a ciliated graph, Fock-Rosly attach a Poisson structure on G^{2g+n-1} and shows that the adjoint action of G in this space is Poisson-Lie. Their main result is the following:

Theorem 6.2. *The inherited Poisson structure on G^{2g+n-1}/G does not depend on the choice of the underlying ciliated fat graph, and coincides with the Atiyah–Bott Poisson structure.*

The main result of this section is then:

Theorem 6.3. *The algebra a_P is a quantization of the Fock–Rosly Poisson structure on G^{2g+n-1}. Its invariant part $\mathrm{Hom}_{U_\hbar(\mathfrak{g})}(\mathbb{C}, a_p)$ does not depend on the choice of P and is a quantization of the Atiyah–Bott Poisson structure on S.*

Before the proof, let us reformulate Fock–Rosly's construction in a way convenient for our purpose (we refer the reader to the original paper [FR99] or the survey [Aud97] for details). Fix a ciliated graph Γ with one vertex, let Σ be the corresponding surface and label the edges from 1 to n in the order determined by Γ. Since Γ has only one vertex, the Fock–Rosly Poisson bracket is defined on

$$\mathcal{O}(G)^{\otimes n} = \mathcal{O}(\mathrm{Hom}(\pi_1(\Sigma), G)).$$

Moreover, it is enough to compute the Poisson bracket $\{f, g\}$ for $f \in \mathcal{O}(G)^{(i)}$ and $g \in \mathcal{O}(G)^{(j)}$ for $i \leq j$, which in turn is determined from the Poisson brackets coming from graphs with one vertex and one or two edges.

For $x \in \mathfrak{g}$, we will denote by x^l, x^r the action of x on G by left-invariant and right invariant vector field respectively, and we set $x^{ad} = x^r - x^l$. Let $r \in \mathfrak{g}^{\otimes 2}$ be the classical limit of the quantum R-matrix and set $\rho = \frac{1}{2}(r^{1,2} - r^{2,1})$ and $t = \frac{1}{2}(r^{1,2} + r^{2,1})$. Let π_{STS} be the bivector field

$$\pi_{STS} = \rho^{ad,ad} + t^{r,l} - t^{l,r}.$$

It induces on G a Poisson structure which has been introduced by Semenov–Tian–Shansky [STS94]. We will denote by G_{STS} the variety G equipped with this Poisson structure. Then G_{STS} is a Poisson Lie variety under the adjoint action of the Poisson-Lie group G.

We can then extract the following from [FR99]:

Theorem 6.4. *Let Γ be a ciliated graph with one vertex and n edges. The corresponding Poisson structure on $\mathcal{O}(G)^{\otimes n}$ is induced by the bivector field*

$$\sum_i \pi_{STS}^{(i)} + \sum_{i<j} (\pi_{ij} - \pi_{ji})$$

where it is understood that π_{ij} is a 2-tensor acting on the ith component of the first factor and the jth component of the second factor of $G^{2n} = G^n \times G^n$, defined by:

$$\pi_{ij} = \begin{cases} \pm(r^{ad,ad}) & \text{if } i,j \text{ are } \pm \text{ unlinked} \\ \pm(r^{ad,ad} + 2t^{r,l}) & \text{if } i,j \text{ are } \pm \text{ linked} \\ \pm(r^{ad,ad} - 2t^{r,r} + 2t^{r,l}) & \text{if } i,j \text{ are } \pm \text{ nested} \end{cases}$$

6.3. Quantization via $\mathrm{Rep}_\hbar G$.

Proof of Theorem 6.3. The following is well known (see e.g. [DM03, Mud06]):

Theorem 6.5. *The reflection equation algebra $\mathcal{O}(\mathrm{Rep}_\hbar G)$ is a flat deformation of $\mathcal{O}(G)$, quantizing G_{STS}.*

Moreover, the left $U_\hbar(\mathfrak{g})^{\otimes 2}$ action on the reflection equation algebra becomes, at the classical limit, the left $U(\mathfrak{g})^{\otimes 2}$ action on $\mathcal{O}(G)$, where the action of the first copy of $U(\mathfrak{g})$ is induced by

$$\mathfrak{g} \ni x \mapsto x^r$$

and the second action is induced by

$$\mathfrak{g} \ni x \mapsto -x^l$$

(since left invariant vector fields act on the right, the minus sign turns this into a left action). Hence to prove the claim it is enough to check that say for $i < j$ the bivector field π_{ij} coincides with the quasi-classical limit of the action of L_{ij}, N_{ij} or U_{ij} if the handles i, j are linked, nested or unlinked respectively. We prove this in the nested case, the other cases are similar: dropping indices for the sake of clarity, we have

$$N = \tau_{12,34} \circ R_{1,34}(R^{-1})_{34,2} = \tau_{12,34} \circ R_{12,34}(R_{34,2}R_{2,34})^{-1}, \tag{6.1}$$

and hence,

$$\frac{\tau_{12,34} \circ N - 1}{\hbar} \mod (\hbar) = r^{12,34} - 2t^{1,3} - 2t^{1,4}. \tag{6.2}$$

where we used that $r^{i,j} + r^{j,i} = 2t^{i,j}$ and

$$r^{ij,k} = r^{i,k} + r^{j,k} \qquad\qquad r^{i,jk} = r^{i,j} + r^{i,k}.$$

The right hand side of (6.2) acts on $G \times G$ via the bivector field

$$r^{ad,ad} - 2t^{r,r} + 2t^{r,l}$$

as required. ☐

References

[AB83] M. F. Atiyah, R. Bott. The Yang-Mills equations over Riemann surfaces. *Philos. Trans. Roy. Soc. London Ser. A* (**1983**). 308(1505):523–615.

[AF12] D. Ayala, J. Francis. Factorization homology of topological manifolds. *ArXiv:1206.5522* (**2012**).

[AFT14] D. Ayala, J. Francis, H. Tanaka. Factorization homology of stratified spaces. *ArXiv:1409.0848* (**2014**).

[AG14] D. Arinkin, D. Gaitsgory. Singular support of coherent sheaves and the geometric Langlands conjecture. *Selecta Mathematica* (**2014**). pp. 1–199.

[AGS96] A. Alekseev, H. Grosse, V. Schomerus. Combinatorial quantization of the Hamiltonian Chern-Simons theory II. *Communications in Mathematical Physics* (**1996**). 174(3):561–604.

[AKSM02] A. Alekseev, Y. Kosmann-Schwarzbach, E. Meinrenken. Quasi-Poisson manifolds. *Canad. J. Math.* (**2002**). 54(1):3–29.

[Ale93] A. Alekseev. Integrability in Hamiltonian Chern-Simons theory. *ArXiv:hep-th/9311074* (**1993**).

[AM95] A. Alekseev, A. Malkin. Symplectic structure of the moduli space of flat connection on a Riemann surface. *Communications in Mathematical Physics* (**1995**). 169(1):99–119.

[APK91] H. H. Andersen, P. Polo, W. Kexin. Representations of quantum algebras. *Inventiones mathematicae* (**1991**). 104(1):1–59.

[AS96] A. Y. Alekseev, V. Schomerus. Representation theory of Chern-Simons observables. *Duke Math. J.* (**1996**). 85(2):447–510.

[Aud97] M. Audin. Lectures on gauge theory and integrable systems. In J. Hurtubise, F. Lalonde, G. Sabidussi, editors, *Gauge Theory and Symplectic Geometry*, vol. 488 of *NATO ASI Series*, pp. 1–48 (Springer Netherlands, **1997**).

[BD04] A. Beilinson, V. Drinfeld. *Chiral algebras*, vol. 51 of *American Mathematical Society Colloquium Publications* (American Mathematical Society, Providence, RI, **2004**).

[BJ14] A. Brochier, D. Jordan. Fourier transform for quantum D-modules via the punctured torus mapping class group. *ArXiv:1403.1841* (**2014**).

[BK06] E. Backelin, K. Kremnizer. Quantum flag varieties, equivariant quantum D-modules, and localization of quantum groups. *Advances in Mathematics* (**2006**). 203(2):408 – 429.

[BK08] E. Backelin, K. Kremnizer. Localization for quantum groups at a root of unity. *J. Amer. Math. Soc.* (**2008**). 21(4):1001–1018.

[BKP89] R. Blackwell, G. M. Kelly, A. J. Power. Two-dimensional monad theory. *J. Pure Appl. Algebra* (**1989**). 59(1):1–41.

[BN11] A. Bruguires, S. Natale. Exact sequences of tensor categories. *International Mathematics Research Notices* (**2011**). 2011(24):5644–5705.

[BZBJ16] D. Ben-Zvi, A. Brochier, D. Jordan. Quantum character varieties and braided module categories. *ArXiv:1606.04769* (**2016**).

[BZFN10] D. Ben-Zvi, J. Francis, D. Nadler. Integral transforms and Drinfeld centers in derived algebraic geometry. *J. Amer. Math. Soc.* (**2010**). 23(4):909–966.

[BZN09] D. Ben-Zvi, D. Nadler. The character theory of a complex group. *arXiv preprint arXiv:0904.1247* (**2009**).

[BZN16] D. Ben-Zvi, D. Nadler. Betti geometric Langlands. *In preparation* (**2016**).

[CEE09] D. Calaque, B. Enriquez, P. Etingof. Universal KZB equations: The elliptic case. In Y. Tschinkel, Y. Zarhin, editors, *Algebra, Arithmetic, and Geometry*, vol. 269 of *Progress in Mathematics*, pp. 165–266 (Birkhuser Boston, **2009**).

[CG] K. Costello, O. Gwilliam. *Factorization algebras in perturbative quantum field theory* (Preprint, Available at `http://www.math.northwestern.edu/~costello/`).

[CKY97] L. Crane, L. H. Kauffman, D. N. Yetter. State-sum invariants of 4-manifolds. *J. Knot Theory Ramifications* (**1997**). 6(2):177–234.

[CP94] V. Chari, A. Pressley. *A guide to quantum groups* (Cambridge University Press, Cambridge, **1994**).

[CS] D. Calaque, C. Scheimbauer. Factorization homology as a fully extended topological field theory. *(In preparation)* .

[DCK90] C. De Concini, V. G. Kac. Representations of quantum groups at roots of 1. In *Operator algebras, unitary representations, enveloping algebras, and invariant theory (Paris, 1989)*, vol. 92 of *Progr. Math.*, pp. 471–506 (Birkhäuser Boston, Boston, MA, **1990**).

[DKM03] J. Donin, P. P. Kulish, A. I. Mudrov. On a universal solution to the reflection equation. *Lett. Math. Phys.* (**2003**). 63(3):179–194.

[DM02a] J. Donin, A. Mudrov. Explicit equivariant quantization on coadjoint orbits of $GL(n, \mathbb{C})$. *Letters in Mathematical Physics* (**2002**). 62(1):17–32.

[DM02b] J. Donin, A. Mudrov. Reflection equation- and FRT-type algebras. *Czechoslovak J. Phys.* (**2002**). 52(11):1201–1206. Quantum groups and integrable systems (Prague, 2002).

[DM03] J. Donin, A. Mudrov. Reflection equation, twist, and equivariant quantization. *Israel J. Math.* (**2003**). 136:11–28.

[DSPS13] C. L. Douglas, C. Schommer-Pries, N. Snyder. Dualizable tensor categories. *ArXiv:1312.7188* (**2013**).

[DSPS14] C. L. Douglas, C. Schommer-Pries, N. Snyder. The balanced tensor product of module categories. *arXiv preprint arXiv:1406.4204* (**2014**).

[EGNO15] P. Etingof, S. Gelaki, D. Nikshych, V. Ostrik. *Tensor categories*, vol. 205 of *Mathematical Surveys and Monographs* (American Mathematical Society, Providence, RI, **2015**).

[ENO10] P. Etingof, D. Nikshych, V. Ostrik. Fusion categories and homotopy theory. *Quantum Topol.* (**2010**). 1(3):209–273. With an appendix by Ehud Meir.

[Enr14] B. Enriquez. Flat connections on configuration spaces and braid groups of surfaces. *Adv. Math.* (**2014**). 252:204–226.

[EO04] P. Etingof, V. Ostrik. Finite tensor categories. *Mosc. Math. J.* (**2004**). 4(3):627–654.

[Fad00] L. Faddeev. Modular double of a quantum group. In *Conférence Moshé Flato Vol. I*, vol. 21 of *Math. Phys. Stud.* (Kluwer Acad. Publ., Dordrecht, **2000**) pp. 149–156.

[FG00] C. Frohman, R. Gelca. Skein modules and the noncommutative torus. *Transactions of the American Mathematical Society* (**2000**). 352(10):4877–4888.

[FG09] V. Fock, A. Goncharov. The quantum dilogarithm and representations of quantum cluster varieties. *Invent. Math.* (**2009**). 175(2):223–286.

[FI14] I. Frenkel, I. Ip. Positive representations of split real quantum groups and future perspectives. *Int. Math. Res. Not.* (**2014**). (8):2126–2164.

[FR99] V. V. Fock, A. A. Rosly. Poisson structure on moduli of flat connections on Riemann surfaces and the r-matrix. In *Moscow Seminar in Mathematical Physics*, vol. 191 of *Amer. Math. Soc. Transl. Ser. 2*, pp. 67–86 (Amer. Math. Soc., Providence, RI, **1999**).

[FRT90] L. Faddeev, N. Y. Reshetikhin, L. Takhtajan. Quantization of Lie groups and Lie algebras. *Yang-Baxter Equation In Integrable Systems. Series: Advanced Series in Mathematical Physics,* *ISBN: 978-981-02-0120-3. WORLD SCIENTIFIC, Edited by Michio Jimbo, vol. 10, pp. 299-309* (**1990**). 10:299–309.

[FT12] D. S. Freed, C. Teleman. Relative quantum field theory. *ArXiv:1212.1692* (**2012**).

[Gai15] D. Gaitsgory. Sheaves of categories and the notion of 1-affineness. In *Stacks and categories in geometry, topology, and algebra*, vol. 643 of *Contemp. Math.*, pp. 127–225 (Amer. Math. Soc., Providence, RI, **2015**).

[Gin14] G. Ginot. Notes on factorization algebras, factorization homology and applications. *ArXiv:1307.5213* (**2014**).

[Gol84] W. Goldman. The symplectic nature of fundamental groups of surfaces. *Advances in Mathematics* (**1984**). 54(2):200 – 225.

[Hau14] R. Haugseng. The higher Morita category of E_n-algebras. *arXiv preprint arXiv:1412.8459* (**2014**).

[JFS15] T. Johnson-Freyd, C. Scheimbauer. The (op) lax square of an (∞, n)-category. *arXiv preprint arXiv:1502.06526* (**2015**).

[Jor09] D. Jordan. Quantum D-modules, elliptic braid groups, and double affine Hecke algebras. *International Mathematics Research Notices* (**2009**). 2009:2081–2105(25).

[Jor14] D. Jordan. Quantized multiplicative quiver varieties. *Adv. Math.* (**2014**). 250:420–466.

[Kel89] G. M. Kelly. Elementary observations on 2-categorical limits. *Bull. Austral. Math. Soc.* (**1989**). 39(2):301–317.

[KW07] A. Kapustin, E. Witten. Electric-magnetic duality and the geometric Langlands program. *Commun. Number Theory Phys.* (**2007**). 1(1):1–236.

[LBS13] D. Li-Bland, P. Severa. Moduli spaces for quilted surfaces and Poisson structures. *ArXiv:1212.2097* (**2013**).

[LBS14] D. Li-Bland, P. Severa. On deformation quantization of Poisson-Lie groups and moduli spaces of flat connections. *ArXiv:1307.2047* (**2014**).

[LF12] I. Lopez Franco. Tensor products of finitely cocomplete and abelian categories. *ArXiv:1212.1545* (**2012**).

[LN15] S. Lentner, D. Nett. New R-matrices for small quantum groups. *Algebras and Representation Theory* (**2015**). 18(6):1649–1673.

[Lur] J. Lurie. *Higher algebra* (Preprint, Available at http://www.math.harvard.edu/~lurie/).

[Lur09] J. Lurie. On the Classification of Topological Field Theories. *ArXiv:0905.0465* (**2009**).

[Lur16] J. Lurie. *Spectral Algebraic Geometry* (Preprint, Available at http://www.math.harvard.edu/~lurie/, **2016**).

[Lus90] G. Lusztig. Finite dimensional Hopf algebras arising from quantized universal enveloping algebras. *Journal of the American Mathematical Society* (**1990**). 3(1):257–296.

[Mud06] A. Mudrov. On quantization of the Semenov-Tian-Shansky Poisson bracket on simple algebraic groups. *Algebra i Analiz* (**2006**). 18(5):156–172.

[Ost03] V. Ostrik. Module categories, weak Hopf algebras and modular invariants. *Transformation Groups* (**2003**). 8(2):177–206.

[PTVV11] T. Pantev, B. Toën, M. Vaquie, G. Vezzosi. Shifted Symplectic Structures. *arXiv:1111.3209* (**2011**).

[RS02] P. Roche, A. Szenes. Trace functionals on noncommutative deformations of moduli spaces of flat connections. *Advances in Mathematics* (**2002**). 168(2):133 – 192.

[RT90] N. Y. Reshetikhin, V. G. Turaev. Ribbon graphs and their invariants derived from quantum groups. *Comm. Math. Phys.* (**1990**). 127(1):1–26.

[RT91] N. Reshetikhin, V. G. Turaev. Invariants of 3-manifolds via link polynomials and quantum groups. *Invent. Math.* (**1991**). 103(3):547–597.

[Sel11] P. Selinger. A survey of graphical languages for monoidal categories. In B. Coecke, editor, *New Structures for Physics*, vol. 813 of *Lecture Notes in Physics*, pp. 289–355 (Springer Berlin Heidelberg, **2011**).

[STS94] M. A. Semenov-Tian-Shansky. Poisson Lie groups, quantum duality principle, and the quantum double. In *Mathematical aspects of conformal and topological field theories and quantum* *groups (South Hadley, MA, 1992)*, vol. 175 of *Contemp. Math.*, pp. 219–248 (Amer. Math. Soc., Providence, RI, **1994**).

[Toë14] B. Toën. Derived algebraic geometry and deformation quantization. *ArXiv:1403.6995* (**2014**).

[TV14] J. Teschner, G. Vartanov. 6j symbols for the modular double, quantum hyperbolic geometry, and supersymmetric gauge theories. *Lett. Math. Phys* (**2014**). 104(5):527–551.

[Vor99] A. A. Voronov. The Swiss-cheese operad. In *Homotopy invariant algebraic structures (Baltimore, MD, 1998)*, vol. 239 of *Contemp. Math.*, pp. 365–373 (Amer. Math. Soc., Providence, RI, **1999**).

[VV10] M. Varagnolo, E. Vasserot. Double affine Hecke algebras at roots of unity. *Represent. Theory* (**2010**). 14:510–600.

[Wal] K. Walker. *Note on topological field theories* (Available at http://canyon23.net/math/tc.pdf).

Department of Mathematics, University of Texas, Austin, TX 78712-0257
E-mail address: `benzvi@math.utexas.edu`

MPIM, Bonn
E-mail address: `abrochier@mpim-bonn.mpg.de`

School of Mathematics, University of Edinburgh, Edinburgh, UK
E-mail address: `D.Jordan@ed.ac.uk`